



Patient-Reported Outcomes Measurement Information System
Dynamic Tools to Measure Health Outcomes From the Patient Perspective

PROMIS Sexual Function and Satisfaction

Manual



Table of Contents

Introduction	4
Sexual Function and Satisfaction (SexFS) Version 2.0	5
Introductory Video	6
About Sexual Function and Satisfaction	6
Introduction to Assessment Options	7
Version Differences	8
Instrument Differences	8
Scoring the Instrument	9
Validity	11
Preview of Sample Item	11
Frequently Asked Questions (FAQs)	12
Appendix A-Scoring Tables FOR SEXSF V2.0	14
Brief Profile (Female)	14
Brief Profile (Female)	15
Brief Profile (Female)	16
Brief Profile (Male)	17
Brief Profile (Male)	18
Full Profile (Female)	19
Full Profile (Female)	20
Full Profile (Female)	21
Full Profile (Female)	22
Full Profile (Female)	23
Full Profile (Female)	24
Full Profile (Male)	25
Full Profile (Male)	26
Full Profile (Male)	27
Full Profile (Male)	28
Full Profile (Male)	29



Section 1: The PROMIS Sexual Function and Satisfaction Measures Brief Profile v1.0 30

Overview 30

Subdomain Definitions..... 30

Reliability and Validity..... 30

Scoring..... 31

Section 2: Creation of a Customized Sexual Function and Satisfaction Assessment 32

Introduction 32

Instrument Descriptions..... 32

Available Instruments 32

Subdomain Definitions..... 33

Selecting Appropriate Items..... 34

A Note on Response Options for Sexual Activities..... 34

Procedures and Data in Support of Validity and Reliability 35

Procedures for Selecting the PROMIS Sexual Function and Satisfaction Profile in Assessment Center..... 38

Procedures for Creating a Custom PROMIS SexFS Instrument in Assessment Center 39

Procedures for Previewing PROMIS Studies in Assessment Center 39

Contact Us 40

References..... 40

Appendix B - Sexual Function and Satisfaction Measures Brief Profile Look-up Tables 42

Appendix C - Sexual Function and Satisfaction Bank Look-up Tables..... 43

Appendix D. Characteristics of Validation Sample 44



Introduction

This document describes resources for the multiple versions of PROMIS I Function and Satisfaction instruments. It is divided into two parts: the first is for v2.0 instruments and the second is for v1.0 instruments.



Sexual Function and Satisfaction (SexFS) Version 2.0

A brief guide to the PROMIS® SexFS v2.0 domains (Table 1) and profile measures (Table 2)

Table 1. PROMIS SexFS v2.0 Domains

	# of items	Calibrated or Uncalibrated ^a	Scored ^b	Requires Activity Screener? ^c
Satisfaction with Sex Life	5	Calibrated	Scored	Yes
Vaginal Lubrication for Sexual Activity	6	Calibrated	Scored	Yes
Vaginal Discomfort with Sexual Activity	11	Calibrated	Scored	Yes
Erectile Function	11	Calibrated	Scored	Yes
Vulvar Discomfort with Sexual Activity – Labial	4	Calibrated	Scored	Yes
Vulvar Discomfort with Sexual Activity – Clitoral	4	Calibrated	Scored	Yes
Oral Discomfort with Sexual Activity	6	Calibrated	Scored	Yes
Oral Dryness with Sexual Activity	3	Calibrated	Scored	Yes
Orgasm – Pleasure	3	Calibrated	Scored	Yes
Orgasm – Ability	1	Uncalibrated	Scored	Yes
Interest in Sexual Activity	2	Uncalibrated	Scored	No
Bother Regarding Sexual Function	11	Uncalibrated	Unscored	No
Factors Interfering with Sexual Satisfaction	35	Uncalibrated	Unscored	No
Therapeutic Aids for Sexual Activity	7	Uncalibrated	Unscored	No
Sexual Activities	15	Uncalibrated	Unscored	No
Anal Discomfort with Sexual Activity	6	Uncalibrated	Scored	Yes
Sexual Function Screener Items	5	Screener	Unscored	No

^aRefers to whether the domain consists of IRT-calibrated items or not. ^bIndicates whether T-scores centered around U.S. population means are available. ^cIndicates whether domain is only intended for sexual active adults.



Table 2. PROMIS SexFS v2.0 Profiles

	Full Profile Items (#)		Brief Profile Items (#)	
	Sexual activity in past 30 days	No sexual activity in past 30 days	Sexual activity in past 30 days	No sexual activity in past 30 days
Interest in Sexual Activity	2	2	2	2
Sexual Activity Screener	1	1	1	1
Reasons for No Sexual Activity	--	1	--	1
Erectile Function (men)	3	--	2	--
Lubrication (women)	3	--	2	--
Vaginal Discomfort (women)	2	--	2	--
Labial Discomfort (women)	2	--	1	--
Clitoral Discomfort (women)	2	--	1	--
Orgasm Ability	1	--	1	--
Orgasm Pleasure	2	--	1	--
Satisfaction with Sex Life	4	--	2	--
Oral Discomfort	2	--	--	--
Oral Dryness	2	--	--	--
Anal Discomfort	2	--	--	--
Total # of Items (Men)	19	4	9	4
Total # of Domains/scores (Men)	8	1	5	1
Total # of Items (Women)	25	4	13	4
Total # of Domains/scores (Women)	11	1	8	1

Introductory Video

An introductory video about the PROMIS SexFS v2.0 can be found here: https://www.youtube.com/watch?v=CQwo_2GlaJA

About Sexual Function and Satisfaction

The PROMIS SexFS v2.0 measures a range of sexual activities, symptoms, functioning, and evaluation of experiences over the past 30 days. Also included are general screener items that ask about sexual activity in the past 30 days and reasons for not having sexual activity. The SexFS v2.0 measures are universal rather than disease-specific. The SexFS v2.0 includes the following domains:

Screener Items ask about sexual activity in the past 30 days and reasons for not having sexual activity.

Interest in Sexual Activity assesses a conscious awareness of wanting to engage in sexual activity during the past 30 days. Items are gender-nonspecific. Higher scores indicate greater interest.

Satisfaction with Sex Life assesses how satisfying and pleasurable the person regards his or her sexual life in the past 30 days with no limitation on how the person defines “sex life.” Items are gender-nonspecific. Higher scores indicate more satisfying sexual activities.

Orgasm – Ability assesses the ease with which a person has been able to have an orgasm over the past 30 days. It is measured with a single, gender-nonspecific item for which higher scores indicate a greater ability to have an orgasm.



Orgasm – Pleasure assesses how pleasurable or satisfying the person's orgasms have felt in the past 30 days. Items are gender-nonspecific. Higher scores indicate more pleasurable orgasms.

Erectile Function assesses the frequency and quality of achieving and maintaining an erection for sexual activity over the past 30 days. Higher scores indicate better function.

Vaginal Discomfort with Sexual Activity assesses physical discomfort of the vagina during and immediately following sexual activity, including sensations of pain, rubbing, burning, pulling, or ripping experienced over the past 30 days. Higher scores indicate greater discomfort.

Vulvar Discomfort with Sexual Activity – Clitoral assesses the degree of physical discomfort, including pain, of the clitoris experienced with sexual activity in the past 30 days. Higher scores indicate greater discomfort.

Vulvar Discomfort with Sexual Activity – Labial assesses the degree of physical discomfort, including pain, of the labia experienced with sexual activity in the past 30 days. Higher scores indicate greater discomfort.

Vaginal Lubrication for Sexual Activity scale assesses the wetness or dryness of the vagina experienced for sexual activity over the past 30 days. Higher scores indicate greater lubrication.

Anal Discomfort with Sexual Activity assesses physical discomfort, irritation, pain, and/or bleeding around the anus or rectum during or after sexual activity over the past 30 days. Items are gender-nonspecific.

Oral Discomfort with Sexual Activity assesses the degree of physical discomfort in the mouth, including pain and/or irritation, experienced with sexual activity in the past 30 days. Items are gender-nonspecific. Higher scores indicate greater discomfort.

Oral Dryness with Sexual Activity assesses the lack of saliva in the mouth experienced with sexual activity in the past 30 days. Items are gender-nonspecific. Higher scores indicate greater dryness.

Factors Interfering with Sexual Satisfaction is a collection of items, each of which assesses the person's perception of the degree to which various factors affected their satisfaction with sex life in the past 30 days. These factors include symptoms of disease, side effects from treatment, and other experiences that have been identified by patients. Some items are gender-nonspecific. These items are intended to be stand-alone items and do not comprise a unidimensional scale.

Therapeutic Aids for Sexual Activity is a collection of items, each of which assesses the use in the past 30 days of hormones, personal lubrications, medications, or devices intended to allow for or improve sexual function. Some items are gender-nonspecific. These items are intended to be stand-alone items and do not comprise a unidimensional scale.

Sexual Activities is a collection of items, each of which assesses the frequency of engaging in specific affectionate or sexual behaviors either alone or with a partner in the past 30 days. Some items are gender-nonspecific. These items are intended to be "stand-alone" items and do not comprise a unidimensional scale.

Bother Regarding Sexual Function is a collection of items, each of which assesses the degree of bother people report for each specific aspect of sexual functioning in the past 30 days. Some items are gender-nonspecific. These items are intended to be "stand-alone" items and do not comprise a unidimensional scale.

Introduction to Assessment Options

All items in the PROMIS SexFS were not intended to be administered together, as some domains might not be relevant for particular situations and reliable scores can be generated without having to administer all of the items in a domain. Researchers should select the sexual function and satisfaction domains and items that are

relevant to their specific sample. There are multiple assessment options: the full profile, brief profile, and customized short forms of any individual domain (that is, a selection of items from the desired domain). When administering any of these options, instruct participants to answer all of the items (i.e., questions or statements) presented. The tables on the first page show how many items are in each individual domain and profile. Any of these options can be administered on paper or by computer.

Whether one uses a customized short form or profile, for those domains marked “calibrated,” the score metric is derived from Item Response Theory (IRT), a family of statistical models that link individual questions to a presumed underlying trait or concept within the overall domain of sexual function, represented by items in the item banks. When choosing among the options available, it is useful to consider the precision gained with longer instruments versus the psychological, physical, and cognitive burden placed on respondents as a result of the number of questions asked.

Version Differences

Some PROMIS domains have multiple versions of instruments (i.e., v1.0, v1.1, v2.0). Generally, **it is recommended that you use the most recent version available which can be identified as the instrument with the highest version number.**

Building on v1.0 of the PROMIS Sexual Function and Satisfaction measure (SexFS), a comprehensive development process was undertaken to create an expanded and improved PROMIS SexFS v2.0. This tool assesses multiple components of sexual functioning, and the validation process included a strong focus on inclusiveness with regard to literacy level, race, age, sexual orientation, and health conditions. Note that while v1.0 was developed exclusively in cancer patients, v2.0 retains the content validity for cancer patients and expands on it, making v2.0 a better measure for cancer populations than v1.0. Scores from v1.0 measures should not be compared to scores from v2.0 measures. Those who have administered items from the SexFS v1.0 and wish to use v2.0 scoring should contact the developers for more information.

Several features of SexFS v2.0 are consistent with v1.0. The measurement system is modular and customizable in that users need only measure those sexual function domains of relevance to their particular study or sample. One important change from v1.0 to 2.0 is that scores are now expressed using a meaningful metric, with scores centered around norms for the population of sexually active US adults. Norms are provided herein for age and sex to aid in the interpretation of SexFS scores. Also, differential item functioning (DIF) was examined by gender, sexual activity, and age to assess the appropriateness of items across different groups of people. Finally, the v2.0 domains demonstrated good convergent and known groups validity and reliability. The final set of items is applicable for both men and women, those sexually active with a partner and without, and those who identify as heterosexual or straight, lesbian, gay, or bisexual.

Instrument Differences

The tables at the beginning of this Manual provide a helpful overview of the different instruments and assessment options. To summarize the tables, of the PROMIS SexFS instruments, 9 are calibrated scales. This means that if one or more items from within that instrument are administered, a respondent’s score will be calculated using item response theory statistics. Three of the instruments are scored but not calibrated. The items within those instruments are combined to create a score, but this score is not based on item response

theory statistics. Five of the instruments do not have calibrated items and are not scored, that is, items within those instruments are not combined in any way to create a score. Each item in these instruments measures a very specific construct corresponding only to that item (e.g., how much radiation burns have affected one's satisfaction with their sex life). For any given item in these uncalibrated instruments, the researcher can use the raw item responses directly for analyses.

Profiles

Profiles are of collections of short forms and items that assess a person with respect to multiple aspects of sexual function and satisfaction. The SexFS currently includes Brief Profiles and Full Profiles (see below). Items were selected for inclusion based on rankings using psychometric and pragmatic criteria: (1) maximum interval information (where applicable, i.e., IRT information curves reflected the greatest degree of precision for the greatest range of the latent variable of interest), (2) consistent item formatting, and (3) content coverage.

Brief Profiles (Male, Female). The Brief Profiles—one for males, one for females—efficiently assess interest in sexual activity followed by a screener item about sexual activity. For those who have not been sexually active in the past 30 days, a fourth and final question asks about reasons for not having sexual activity. For those who have been sexually active in the past 30 days, men are asked about erectile function, while women are asked about vaginal lubrication, vaginal discomfort, and vulvar discomfort; men and women are asked about orgasm and satisfaction with sex life. These are the domains that are most likely relevant for the majority of healthy people and those suffering from chronic diseases.

Full Profiles (Male, Female). The Full Profiles—one for males, one for females—includes all of the domains measured in the Brief Profiles, but also adds oral dryness, oral discomfort, and anal discomfort. For the domains that overlap between the Brief and Full Profiles, the Full Profiles use equal or greater number of items to assess each domain.

Customized Short Forms

Within any of the 9 calibrated scales, users can select one or more items to create a customized short form for measuring that domain. Selection of the items could be based on suitability of the item for the particular population of interest. Note that the items that generally perform the best from the 9 calibrated scales are found on the Brief Profiles and Full Profiles (see above).

Selecting an instrument

In selecting among options, the differences are domain coverage and instrument length. The reliability and precision of the short forms within a domain is highly similar. If you are working with a sample in which you want the most precise measure, select the longest short form. If you have little room for additional measures but really wanted to capture something as a secondary outcome, select one of the shorter instruments (e.g., brief profile or just 1 or 2 domains of interest).

Scoring the Instrument

PROMIS instruments are scored using item-level calibrations. This means that the most accurate way to score a PROMIS instrument is to use the HealthMeasures Scoring Service (https://www.assessmentcenter.net/ac_scoringervice) or a data collection tool that automatically calculates scores (e.g., Assessment Center, REDCap auto-score). This method of scoring uses responses to each item for each participant. We refer to this as “response pattern scoring.” Response pattern scoring is preferred because it is more accurate than the use of the summed raw score/scale score look up tables included in this manual (for the full profile). Response pattern scoring is especially useful when there are missing data (i.e., a respondent

skipped an item), different groups of participants responded to different items, or you have created a new questionnaire using a subset of questions from a PROMIS item bank. Response pattern scoring can be used for the SexFS Brief Profiles, Full Profiles, and customized short forms, as long as scoring software is used such as those mentioned earlier. A video tutorial provides instructions for using the HealthMeasures Scoring Service to score PROMIS SexFX measures including how to handle screener items and “not applicable” responses (<https://www.youtube.com/watch?v=d11SXxfvKu0>).

For the **Brief Profiles**, an alternative to using specialized scoring software for response pattern scoring is to use the response pattern scoring look-up tables in the Appendix, which allow a user to determine the SexFS T-score for any combination of item responses for each domain.

For the **Full Profiles**, the alternative to using specialized scoring software with response pattern scoring is to use a lookup table based on simple summed raw scores. Lookup tables based on response patterns become too onerous with more than two items in a domain, and so these tables allow one to obtain a SexFS T-score corresponding to the sum of the items within the domain of interest. Note that this approach does not result in as precise a score as the response pattern scoring method. The directions below specify how to use the sumscores lookup tables in the Appendix to score the SexFS Full Profiles.

Create a Summed Raw Score. First, a raw summed score is created for each domain in the profile. However, this raw summed domain score can only be created if ALL items from that domain were answered and NO “not applicable” responses were given. For example, if a respondent only answered one of the two Satisfaction with Sex Life items, a valid Satisfaction with Sex Life score cannot be produced. If a respondent answered the Orgasm – Ability item by selecting “Have not tried to have an orgasm in the past 30 days,” a valid Orgasm - Ability score cannot be produced.

After confirming all items in a given domain were answered without endorsing a “not applicable” response (identified by a score of 0), add up the response scores to all items in that domain. This is the raw summed score for that domain. For example, for Satisfaction with Sex Life, the raw summed score can range from 4 to 20. Some domains, such as Orgasm – Ability and Vulvar Discomfort – Labial, contain only 1 item in the Brief Profile, and so no summing is required. **All questions must be answered in order to produce a valid score using the scoring tables.** If a participant has skipped a question, use the HealthMeasures Scoring Service (https://www.assessmentcenter.net/ac_scoringervice) to generate a final score.

Use the Raw Score/T-Score Look-up Tables. Locate the applicable score conversion table in the Appendix and use this table to translate the domain’s summed raw score into a T-score for each participant. The T-score rescales the raw score into a standardized score with a mean of 50 and a standard deviation (SD) of 10. The standardized T-score is reported as the final score for each participant for each domain. Therefore, a person with a T-score of 40 is one SD below the mean.

As an example, for the Satisfaction with Sex Life domain in full profile, a raw summed score of 6 converts to a T-score of 37.57 with a standard error (SE) of 2.00 (see scoring table in Appendix A). Thus, the 95% confidence interval around the observed score ranges from 33.63 (T-score - 1.96*SE) to 41.49 (T-score + 1.96*SE) -.

Scoring When the Respondent Has Not Had Sexual Activity. For many domains (see Table 1), a screener item should be used first that asks whether they have been sexually active in the past 30 days. If they answer “Yes,” then they are administered that domain. If they answer “No,” they are not administered any domains that are intended to provide assessments of functioning with or during sexual activity. However, people who respond that they have not had sexual activity in the past 30 days can be asked an additional item to indicate the reasons they have not had activity using a checklist. This checklist could be used by researchers to differentiate among

respondents who did not have activity due to a sexual concern of interest (e.g., difficulty getting erections) versus those who did not have activity for other reasons (e.g., partner not available).

Scoring For Uncalibrated SexFS Domains

Domains without calibrations are either (1) collections of individual items measuring separate things (e.g., Therapeutic Aids for Sexual Activity) or (2) sets of items that might measure an underlying latent concept, but for which an IRT model was not possible due to small number of items (e.g., Interest in Sexual Activity) or a small number of respondents (e.g., Anal Discomfort with Sexual Activity). We describe each below.

Scoring Uncalibrated Measures of Latent Domains: Interest in Sexual Activity and Orgasm-Ability. For these two domains, there were too few items to use Item Response Theory. Using the look-up tables in the Appendix, for Interest, convert the summed score to a T-score; for Orgasm-Ability, convert the raw score to a T-score.

Scoring Uncalibrated Individual Items: Factors Interfering with Sexual Satisfaction, Therapeutic Aids for Sexual Activity, Sexual Activities). These domains are collections of individual items that each measure an individual concept. Accordingly, the raw responses to the individual item should be taken as the score for that item.

Special Instructions for Scoring Anal Discomfort with Sexual Activity. As described in Weinfurt et al. 2015, until additional data can be collected from samples with varying degrees of anal discomfort, “the PROMIS SexFS working group suggests that 4 different scores are possible: (1) an Anal Discomfort and Pain with Sexual Activity score that is the average of the 4 items measuring discomfort and pain, (2) an Anal Bleeding with Sexual Activity score that is the average of the 2 items measuring bleeding, (3) a composite score (average of all 6 items) with greater weight given to discomfort/pain than bleeding (Cronbach’s alpha = 0.93), and a composite score that gives equal weight to discomfort/pain and bleeding by averaging (1) and (2).”

Validity

Multiple types of validity have been evaluated and results are described in the main development paper:

Weinfurt KP, Lin L, Bruner DW, Cyranowski JM, Dombek CB, Hahn EA, et al. Development and Initial Validation of the PROMIS((R)) Sexual Function and Satisfaction Measures Version 2.0. *J Sex Med.* 2015;12(9):1961-74. doi: 10.1111/jsm.12966. PubMed PMID: 26346418.

Preview of Sample Item

Below is an excerpt from the paper version of the Brief Profile (Female). This is the paper version format used for all profile instruments. It is important to note, automatic scoring is not available for paper administration.

SFSAT101r	In the past 30 days , how satisfied have you been with your sex life?
	<input type="checkbox"/> 1 Not at all
	<input type="checkbox"/> 2 A little bit
	<input type="checkbox"/> 3 Somewhat
	<input type="checkbox"/> 4 Quite a bit
	<input type="checkbox"/> 5 Very
SFSAT102r	In the past 30 days , how much pleasure has your sex life given you?
	<input type="checkbox"/> 1 None
	<input type="checkbox"/> 2 A little bit
	<input type="checkbox"/> 3 Some
	<input type="checkbox"/> 4 Quite a bit
	<input type="checkbox"/> 5 A lot

Frequently Asked Questions (FAQs)

Q: I am interested in learning more. Where can I do that?

Review the HealthMeasures website at www.healthmeasures.net.

Q: Do I need to register with PROMIS to use these instruments?

No.

Q: Is the SexFS appropriate for LGBT populations?

Self-identified lesbian, gay, and bisexual individuals were engaged in the development process in focus groups, cognitive interviews, and item testing, and item wording is generally appropriate regardless of sexual orientation. However, transgender participants were not a particular focus for the v1.0 or v2.0 measure validation efforts; additional work will be needed to demonstrate validity for transgender participants.

Q: I'm worried that patients might be too embarrassed or offended to answer some of these questions. What has been the experience of PROMIS with this measure?

In testing v1.0 of this measure in a large population of patients with cancer, there were no more missing data on the sexual function measure than other measures on different topics that were being tested simultaneously (e.g., sleep quality, cognitive function, and illness impact). Since then, v2.0 was tested in additional populations and settings and missing data have not been a significant problem. Skipped items can be minimized by informing study participants at the outset that your study will be asking about different domains of health that are important to understand, including (for example) physical function, sleep, sexual function, and fatigue.

Q: Are these instruments available in other languages?

Yes! Look at the HealthMeasures website (www.healthmeasures.net) for current information on PROMIS translations.

Q: Can I make my own short form?

Yes, custom short forms can be made by selecting any items from an item bank. This can be scored using the Scoring Service (https://www.assessmentcenter.net/ac_scoringervice).

Q: How do I handle multiple responses when administering a short form on paper?

Guidelines on how to deal with multiple responses have been established. Resolution depends on the responses noted by the research participant.

- If two or more responses are marked by the respondent, and they are next to one another, then a data entry specialist will be responsible for randomly selecting one of them to be entered and will write down on the form which answer was selected. Note: To randomly select one of two responses, the data entry specialist will flip a coin (heads - higher number will be entered; tails – lower number will be entered). To randomly select one of three (or more) responses, a table of random numbers should be used with a statistician's assistance.
- If two or more responses are marked, and they are NOT all next to one another, the response will be considered missing.

Q: What is the minimum change on a PROMIS instrument that represents a clinically meaningful difference?

To learn more about research on the meaning of a change in scores, we suggest conducting a literature review to identify the most current information. The HealthMeasures website (<http://www.healthmeasures.net/score-and-interpret/interpret-scores/promis>) has additional information on interpreting scores.

Q: Can I compare adult PROMIS SexFS v1.0 and v2.0 scores if the calibrations changed?

Those who have administered items from the SexFS v1.0 and wish to use v2.0 scoring should contact the developers for more information.

Appendix A-Scoring Tables FOR SEXSF V2.0

Brief Profile (Female)

Interest in Sexual Activity Short Form Conversion Table	
Raw Score	T-Score
2	21.9
3	27.4
4	32.9
5	38.4
6	43.9
7	49.4
8	54.8
9	60.3
10	65.8

Orgasm – Pleasure Short Form Conversion Table		
Raw Score	T-Score	SE ¹
1	27.5	4.3
2	34.1	3.1
3	40.2	3.3
4	47.5	3.5
5	58.6	6.2

¹SE: Standard Error

Note: This domain has not been IRT calibrated because it only has 2 items. For any domain that not calibrated, 1) there is no difference in T-Scores between response patterns 1-2 vs 2-1 etc., so summed score is used instead of pattern; 2) There is no SE associated with T-Score since the T-Score is a simple transformation of the observed average score (centered by the general population weighted mean and SD).

Orgasm – Ability Short Form Conversion Table	
Raw Score	T-Score
1	20.8
2	30.2
3	39.6
4	49.0
5	58.4

Note: This domain has not been IRT calibrated because it only has 1 item. There is no SE associated with the T-Score since the T-Score is a simple transformation of the observed average score (centered by the general population weighted mean and SD).



Brief Profile (Female)

Satisfaction with Sex Life Short Form Conversion Table			
SFSAT101	SFSAT102	T-Score	SE ¹
1	1	31.6	4.5
1	2	37.1	2.5
1	3	39.9	2.8
1	4	43.1	3.8
1	5	46.2	5.3
2	1	37.6	2.5
2	2	39.8	2.1
2	3	42.3	2.3
2	4	45.2	3.0
2	5	48.3	4.3
3	1	40.5	3.1
3	2	42.5	2.5
3	3	45.3	2.4
3	4	48.4	2.5
3	5	51.7	3.3
4	1	44.7	4.2
4	2	46.2	3.3
4	3	48.7	2.6
4	4	51.9	2.5
4	5	55.4	2.6
5	1	48.8	5.5
5	2	50.0	4.6
5	3	52.2	3.5
5	4	55.7	2.7
5	5	62.7	5.2

¹SE: Standard Error

Vaginal Lubrication for Sexual Activity Short Form Conversion Table			
SFLUB004	SFLUB001	T-Score	SE ¹
1	1	27.6	4.5
1	2	32.0	3.8
1	3	34.2	4.2
1	4	35.8	4.9
1	5	37.7	5.9
2	1	33.1	3.3
2	2	35.4	3.1
2	3	37.2	3.4
2	4	38.8	3.9
2	5	40.7	4.9
3	1	36.2	3.3
3	2	38.0	3.0
3	3	39.5	3.0
3	4	41.1	3.4
3	5	43.1	4.3
4	1	40.3	4.0
4	2	41.8	3.5
4	3	43.4	3.3
4	4	45.4	3.4
4	5	48.1	3.8
5	1	47.4	4.9
5	2	48.1	4.5
5	3	49.2	4.2
5	4	51.2	4.0
5	5	58.4	6.5

¹SE: Standard Error



Brief Profile (Female)

Vaginal Discomfort for Sexual Activity Short Form Conversion Table			
SFVAG202	SFVAG206	T-Score	SE ¹
1	1	43.3	6.6
1	2	52.8	2.6
1	3	55.6	3.4
1	4	56.8	4.3
1	5	57.3	4.8
2	1	53.3	2.6
2	2	56.9	2.5
2	3	60.4	2.5
2	4	62.5	3.1
2	5	63.6	3.9
3	1	57.3	3.2
3	2	60.9	2.4
3	3	63.8	2.2
3	4	66.4	2.4
3	5	68.3	3.1
4	1	59.8	4.4
4	2	63.7	3.0
4	3	66.8	2.3
4	4	69.9	2.4
4	5	73.0	2.5
5	1	61.9	5.7
5	2	66.2	4.2
5	3	69.9	3.1
5	4	73.6	2.5
5	5	78.1	3.4

¹SE: Standard Error

Vulvar Discomfort with Sexual Activity – Labial Short Form Conversion Table		
Raw Score	T-Score	SE ¹
1	47.4	8.2
2	62.7	3.2
3	67.9	3.1
4	72.1	3.1
5	77.3	4.0

¹SE: Standard Error

Vulvar Discomfort with Sexual Activity – Clitoral Short Form Conversion Table		
Raw Score	T-Score	SE ¹
1	48.2	8.6
2	65.0	3.4
3	70.4	3.5
4	76.4	4.3

Note: To score the item, responses “Quite a bit” and “A lot” are combined and to be scored as a 4.

Brief Profile (Male)

Interest in Sexual Activity Short Form Conversion Table	
Raw Score	T-Score
2	21.9
3	27.4
4	32.9
5	38.4
6	43.9
7	49.4
8	54.8
9	60.3
10	65.8

Note: This domain has not been IRT calibrated because it only has 2 items. For any domain that not calibrated, 1) there is no difference in T-Scores between response patterns 1-2 vs 2-1 etc., so summed score is used instead of pattern; 2) There is no SE associated with T-Score since the T-Score is a simple transformation of the observed average score (centered by the general population weighted mean and SD).

Orgasm – Ability Short Form Conversion Table	
Raw Score	T-Score
1	20.8
2	30.2
3	39.6
4	49.0
5	58.4

Note: This domain has not been IRT calibrated because it only has 1 item. There is no SE associated with the T-Score since the T-Score is a simple transformation of the observed average score (centered by the general population weighted mean and SD).

Orgasm – Pleasure Short Form Conversion Table		
Raw Score	T-Score	SE ¹
1	27.5	4.3
2	34.1	3.1
3	40.2	3.3
4	47.5	3.5
5	58.6	6.2

¹SE: Standard Error

Erectile Function Short Form Conversion Table			
SFEFN005	SFEFN008	T-Score	SE ¹
1	1	28.7	4.1
1	2	34.3	2.4
1	3	36.6	2.7
1	4	38.8	3.1
1	5	42.6	4.0
2	1	33.5	2.5
2	2	36.2	2.1
2	3	38.2	2.1
2	4	40.1	2.5
2	5	43.5	3.4
3	1	35.5	2.6
3	2	37.7	2.1
3	3	39.4	1.9
3	4	41.2	2.2
3	5	44.3	3.0
4	1	37.0	3.0
4	2	39.2	2.3
4	3	40.8	2.0
4	4	42.7	2.1
4	5	45.8	2.7
5	1	39.2	3.8
5	2	41.3	3.0
5	3	42.8	2.6
5	4	45.2	2.5
5	5	55.7	6.9

¹SE: Standard Error



Brief Profile (Male)

Satisfaction with Sex Life Short Form Conversion Table			
SFSAT101	SFSAT102	T-Score	SE ¹
1	1	31.6	4.5
1	2	37.1	2.5
1	3	39.9	2.8
1	4	43.1	3.8
1	5	46.2	5.3
2	1	37.6	2.5
2	2	39.8	2.1
2	3	42.3	2.3
2	4	45.2	3.0
2	5	48.3	4.3
3	1	40.5	3.1
3	2	42.5	2.5
3	3	45.3	2.4
3	4	48.4	2.5
3	5	51.7	3.3
4	1	44.7	4.2
4	2	46.2	3.3
4	3	48.7	2.6
4	4	51.9	2.5
4	5	55.4	2.6
5	1	48.8	5.5
5	2	50.0	4.6
5	3	52.2	3.5
5	4	55.7	2.7
5	5	62.7	5.2

¹SE: Standard Error



Full Profile (Female)

Interest in Sexual Activity Short Form Conversion Table	
Raw Score	T-Score
2	21.9
3	27.4
4	32.9
5	38.4
6	43.9
7	49.4
8	54.8
9	60.3
10	65.8

Lubrication Short Form Conversion Table		
Raw Score	T-Score	SE
3	25.7	4.1
4	30.2	2.9
5	32.6	2.7
6	34.5	2.7
7	36.2	2.7
8	38.0	2.7
9	39.9	3.0
10	41.9	3.1
11	43.7	3.2
12	45.7	3.2
13	48.3	3.3
14	51.8	3.6
15	59.3	6.2

Vaginal Discomfort Short Form Conversion Table		
Raw Score	T-Score	SE
2	43.1	6.5
3	52.8	2.5
4	56.6	2.4
5	60.3	2.3
6	63.3	2.2
7	66.2	2.2
8	69.3	2.3
9	72.6	2.4
10	77.4	3.5



Full Profile (Female)

Vaginal Discomfort Short Form Conversion Table			
SFVAG202	SFVAG203	T-Score	SE
1	1	43.1	6.5
1	2	52.5	2.5
1	3	55.2	3.3
1	4	56.3	4.2
1	5	56.8	4.7
2	1	53.0	2.4
2	2	56.6	2.4
2	3	60.0	2.4
2	4	62.1	3
2	5	63.2	3.7
3	1	57.2	3.1
3	2	60.5	2.3
3	3	63.3	2.1
3	4	65.9	2.3
3	5	67.8	3
4	1	59.8	4.3
4	2	63.4	2.8
4	3	66.4	2.2
4	4	69.3	2.2
4	5	72.3	2.4
5	1	62.0	5.6
5	2	65.9	4
5	3	69.4	2.9
5	4	72.9	2.3
5	5	77.4	3.4

Vulvar Discomfort – Labial Short Form Conversion Table		
Raw Score	T-Score	SE
2	47.1	8.0
3	60.5	2.3
4	63.7	2.0
5	66.4	2.0
6	68.9	1.9
7	71.1	2.0
8	73.6	2.1
9	77.8	3.5

Vulvar Discomfort – Labial Short Form Conversion Table			
SFVUL203	SFVUL204	T-Score	SE
1	1	47.1	8.0
1	2	61.3	2.1
1	3	64.4	2.8
1	4	66.9	3.8
2	1	60.2	2.3
2	2	63.8	1.9
2	3	66.9	2.0
2	4	69.5	2.6
3	1	62.3	2.6
3	2	66.2	1.9
3	3	69.0	1.8
3	4	71.7	2.0
4	1	63.0	3.0
4	2	67.6	2.3
4	3	70.8	1.8
4	4	73.8	2.0
5	1	63.2	3.2
5	2	68.2	2.7
5	3	72.1	2.2
5	4	77.7	3.4



Full Profile (Female)

Vulvar Discomfort – Clitoral		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	48.0	8.5
3	63.1	2.7
4	66.6	2.2
5	69.4	2.2
6	72.6	2.1
7	75.3	2.3
8	80.0	3.3

Orgasm Ability	
Short Form Conversion Table	
Raw Score	T-Score
1	20.8
2	30.2
3	39.6
4	49.0
5	58.4

Vulvar Discomfort – Clitoral			
Short Form Conversion Table			
SFVUC203	SFVUC204	T-Score	SE
1	1	48.0	8.5
1	2	64.3	2.3
1	3	68.1	3.0
1	4	71.7	4.1
2	1	62.6	2.6
2	2	66.7	2.1
2	3	70.4	2.2
2	4	73.7	2.9
3	1	64.3	2.8
3	2	69.1	2.1
3	3	72.7	2.0
3	4	76.2	2.2
4	1	64.8	3.1
4	2	70.5	2.5
4	3	74.9	2.2
4	4	79.9	3.1



Full Profile (Female)

Orgasm Pleasure		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	25.2	3.9
3	30.1	2.8
4	33.1	2.6
5	36.0	2.7
6	39.3	2.7
7	42.8	2.8
8	47.3	2.9
9	51.8	3.0
10	60.0	5.8

Orgasm Pleasure			
Short Form Conversion Table			
SFOGP201	SFOGP203	T-Score	SE
1	1	25.2	3.9
1	2	30.9	2.6
1	3	35.4	3.2
1	4	41.5	3.8
1	5	48.7	4.5
2	1	29.5	2.7
2	2	33.0	2.4
2	3	36.8	2.7
2	4	42.1	3.4
2	5	48.8	4.3
3	1	32.2	3.1
3	2	35.4	2.5
3	3	39.3	2.6
3	4	43.7	2.8
3	5	49.5	3.7
4	1	34.9	4.2
4	2	38.0	3.2
4	3	42.3	2.7
4	4	47.2	2.9
4	5	52.5	3.0
5	1	37.4	5.9
5	2	40.8	4.7
5	3	45.5	3.6
5	4	50.9	2.9
5	5	60.0	5.8



Full Profile (Female)

Oral Discomfort		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	44.8	7.2
3	55.1	3.6
4	58.6	3.2
5	61.3	3.0
6	63.9	2.9
7	66.7	3.1
8	71.9	4.4

Oral Discomfort			
Short Form Conversion Table			
SFODS201	SFODS202	T-Score	SE
1	1	44.8	7.2
1	2	56.3	3.2
1	3	59.1	3.7
1	4	61.9	4.7
2	1	54.7	3.6
2	2	59.0	2.7
2	3	61.8	2.9
2	4	64.7	3.6
3	1	57.1	3.8
3	2	61.1	2.7
3	3	64.0	2.7
3	4	67.3	3.1
4	1	58.3	4.3
4	2	62.8	3.2
4	3	66.2	2.9
4	4	71.9	4.3



Full Profile (Female)

Anal Discomfort	
Short Form Conversion Table	
Raw Score	T-Score
2	48.6
3	66.0
4	83.5
5	100.9
6	118.3
7	135.7
8	153.2
10	188.0

Satisfaction with Sex Life		
Short Form Conversion Table		
Raw Score	T-Score	SE
3	31.2	4.4
4	36.4	2.5
5	38.7	2.3
6	40.7	2.3
7	42.6	2.3
8	44.5	2.3
9	46.4	2.4
10	48.5	2.4
11	50.7	2.4
12	52.9	2.5
13	55.5	2.6
14	58.8	3.0
15	64.8	5.2



Full Profile (Male)

Interest in Sexual Activity Short Form Conversion Table	
Raw Score	T-Score
2	21.9
3	27.4
4	32.9
5	38.4
6	43.9
7	49.4
8	54.8
9	60.3
10	65.8

IIEF- Erectile Function Short Form Conversion Table		
Raw Score	T-Score	SE
3	28.2	4.0
4	32.6	2.1
5	34.6	1.8
6	36.0	1.8
7	37.1	1.7
8	38.2	1.7
9	39.3	1.7
10	40.4	1.7
11	41.5	1.8
12	42.8	1.8
13	44.3	1.9
14	46.5	2.3
15	56.1	6.8

Orgasm Ability Short Form Conversion Table	
Raw Score	T-Score
1	20.8
2	30.2
3	39.6
4	49.0
5	58.4



Full Profile (Male)

Orgasm Pleasure		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	25.2	3.9
3	30.1	2.8
4	33.1	2.6
5	36.0	2.7
6	39.3	2.7
7	42.8	2.8
8	47.3	2.9
9	51.8	3.0
10	60.0	5.8

Orgasm Pleasure			
Short Form Conversion Table			
SFOGP201	SFOGP203	T-Score	SE
1	1	25.2	3.9
1	2	30.9	2.6
1	3	35.4	3.2
1	4	41.5	3.8
1	5	48.7	4.5
2	1	29.5	2.7
2	2	33.0	2.4
2	3	36.8	2.7
2	4	42.1	3.4
2	5	48.8	4.3
3	1	32.2	3.1
3	2	35.4	2.5
3	3	39.3	2.6
3	4	43.7	2.8
3	5	49.5	3.7
4	1	34.9	4.2
4	2	38.0	3.2
4	3	42.3	2.7
4	4	47.2	2.9
4	5	52.5	3.0
5	1	37.4	5.9
5	2	40.8	4.7
5	3	45.5	3.6
5	4	50.9	2.9
5	5	60.0	5.8



Full Profile (Male)

Oral Discomfort		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	44.8	7.2
3	55.1	3.6
4	58.6	3.2
5	61.3	3.0
6	63.9	2.9
7	66.7	3.1
8	71.9	4.4

Oral Discomfort			
Short Form Conversion Table			
SFODS201	SFODS202	T-Score	SE
1	1	44.8	7.2
1	2	56.3	3.2
1	3	59.1	3.7
1	4	61.9	4.7
2	1	54.7	3.6
2	2	59.0	2.7
2	3	61.8	2.9
2	4	64.7	3.6
3	1	57.1	3.8
3	2	61.1	2.7
3	3	64.0	2.7
3	4	67.3	3.1
4	1	58.3	4.3
4	2	62.8	3.2
4	3	66.2	2.9
4	4	71.9	4.3



Full Profile (Male)

Oral Dryness		
Short Form Conversion Table		
Raw Score	T-Score	SE
2	40.8	6.1
3	49.3	3.3
4	53.2	3.1
5	56.8	3.1
6	60.1	2.9
7	63.4	3.1
8	69.1	4.6

Anal Discomfort	
Short Form Conversion Table	
Raw Score	T-Score
2	48.6
3	66.0
4	83.5
5	100.9
6	118.3
7	135.7
8	153.2
10	188.0

Oral Dryness			
Short Form Conversion Table			
SFODR202	SFODR203	T-Score	SE
1	1	40.8	6.1
1	2	49.9	3.2
1	3	54.6	3.9
1	4	57.7	5.0
2	1	48.5	3.3
2	2	53.3	3.0
2	3	57.5	3.0
2	4	60.5	3.7
3	1	51.2	3.9
3	2	56.5	3.1
3	3	60.3	2.7
3	4	63.8	3.1
4	1	52.7	4.7
4	2	58.9	3.6
4	3	63.0	3.0
4	4	69.1	4.6



Full Profile (Male)

Satisfaction with Sex Life		
Short Form Conversion Table		
Raw Score	T-Score	SE
3	31.2	4.4
4	36.4	2.5
5	38.7	2.3
6	40.7	2.3
7	42.6	2.3
8	44.5	2.3
9	46.4	2.4
10	48.5	2.4
11	50.7	2.4
12	52.9	2.5
13	55.5	2.6
14	58.8	3.0
15	64.8	5.2



Section 1: The PROMIS Sexual Function and Satisfaction Measures Brief Profile v1.0

Overview

The PROMIS Sexual Function and Satisfaction Measures Brief Profile (PSxFBP) provides scores on 7 different subdomains of sexual function: Interest in Sexual Activity, Vaginal Discomfort (women only), Lubrication (women only), Erectile Function (men only), Orgasm, and Global Satisfaction with Sex Life (see below for subdomain definitions). The PSxFBP is intended for broad use, although almost all of the development work was with cancer populations. (Research is ongoing to expand development beyond cancer.) The PSxFBP is available for men and women and consists of the best items selected from each subdomain for general purposes. Each question asks respondents to report on their experiences over the past 30 days.

Subdomain Definitions

Global Satisfaction with Sex Life is the person's overall evaluation of his or her sex life. No limitation is placed on what the person includes in his or her definition of "sex life." Higher scores indicate more satisfaction with sex life. Lower scores indicate less satisfaction with sex life.

Interest in Sexual Activity refers to a conscious awareness of wanting to engage in sexual activity. Items are gender-neutral. Higher scores indicate more interest. Lower scores indicate less interest.

Lubrication refers to the wetness or dryness of the vagina during sexual activity. Higher scores indicate more lubrication. Lower scores indicate less lubrication.

Vaginal Discomfort refers to the degree of physical discomfort of the vagina during and immediately following sexual activity. Higher scores indicate more discomfort as reflected by pain and/or uncomfortable tightness. Lower scores indicate less discomfort as indicated by no pain, bleeding, and/or uncomfortable tightness.

Erectile Function refers to the ability to achieve and maintain an erection for sexual activity. Higher scores indicate better function. Lower scores indicate poorer function.

Orgasm assesses the degree to which the person has experienced a satisfying climax. It is measured with a single, gender-neutral item for which higher scores indicate a greater ability to have satisfying orgasms, and lower scores indicate less ability.

Note: Additional subdomains not included in the PSxFBP are Interfering Factors, Therapeutic Aids, Sexual Activities, and Anal Discomfort. See Section 2 for definitions.

Reliability and Validity

A detailed account of the development of the PROMIS[®] Sexual Function domain, including reliability and validity data, is found in Section 2. Correlations between the PSxFBP and corresponding subdomains of two well-established measures—the FSFI and the IIEF—ranged between .48 and .92 (see Table 3). The subdomains of the PSxFBP discriminate between people who had and had not asked an oncology provider about sexual problems (Table 4). Test-retest correlations are > .65 for all subdomains of the PSxFBP (see Table 5).

Scoring

PROMIS instruments are scored using item-level calibrations. This means that the most accurate way to score a PROMIS instrument is to utilize scoring tools within Assessment Center or API that look at responses to each item for each participant. Data collected in either of these platforms will automatically score in this way. We refer to this as “response pattern scoring.” Response pattern scoring can be used when data was collected on paper or in another software package through the [Assessment Center Scoring Service](#). Because response pattern scoring is more accurate than the use of raw score/scale score look up tables, it is preferred. However, if you aren’t able to use response pattern scoring, you can use the instructions below which rely on raw score/scale score look-up tables.

With the exception of the Orgasm subdomain, all subdomain scores are expressed as *T* scores (mean = 50, standard deviation = 10). At present, a *T* score of 50 corresponds to the mean response among the cancer survivors used for item testing (total *N* = 819). If the PSxFBP is administered electronically using Assessment Center™, scoring is done automatically by the software and scores for every subdomain are added as new variables.

Multiple Domains

The score metric for PROMIS instruments is Item Response Theory (IRT), a family of statistical models that link individual questions to a presumed underlying trait or concept represented by all items in the item bank. In the case of the PROMIS Sexual Function and Satisfaction Brief Profile, the instrument is made up of six individual short forms that are scored individually: Interest in Sexual Activity, Vaginal Discomfort (women only), Lubrication (women only), Erectile Function (men only), Orgasm, and Global Satisfaction with Sex Life. Scoring uses item-level calibrations. This means that the most accurate way to score a PROMIS Profile is to utilize scoring tools within Assessment Center that look at responses to each item for each participant. We refer to this as “response pattern scoring.” Response pattern scoring tools within Assessment Center can be used even if data was collected on paper or in another software package. Because response pattern scoring is more accurate than the use of raw score/scale score look up tables, it is preferred. However, if you aren’t able to use response pattern scoring, you can use the instructions below which rely on raw score/scale score look-up tables (see tables in Appendix B).

Not Applicable Responses

Each question has multiple response options ranging in value from one to four or five. In many cases, there is also a response option that allows a respondent to report a “not applicable” response. For example, for Erectile Function, a respondent can answer “have not tried to get an erection in the past 30 days.” These “not applicable” responses cannot be used to calculate a score. They can only be used as individual items to describe respondents.

Create a Summed Raw Score

A raw summed score is created for each domain in the profile. However, this raw summed domain score can only be created if ALL items from that domain were answered *and* NO “not applicable” responses were given. For example, if a respondent only answered one of the two Global Satisfaction with Sex Life items, a valid Global Satisfaction with Sex Life score cannot be produced. If a different respondent answered both of these items, but endorsed “Have not had sexual activity in the past 30 days” for one or both items, a valid Global Satisfaction with Sex Life score cannot be produced.

After confirming all items in a given domain were answered without endorsing a “not applicable” response (identified by a score of 0), add up the response scores to all items in that domain. This is the raw summed score for that domain. For example, for Global Satisfaction with Sex Life, the raw summed score can range from 2 (endorsed “Not at all” to both items) to 10 (endorsed “Very” or “Very much” to both items).

Note that for the single Orgasm item, no summed score is produced. This item is not scored using Item Response Theory. Instead, raw responses can be used in analyses.

Use the Raw Score/T-Score Look-up Tables

Locate the applicable score conversion table in Appendix B and use this table to translate the domain raw summed score into a T-score for each participant. The T-score rescales the raw score into a standardized score with a mean of 50 and a standard deviation (SD) of 10. Therefore a person with a T-score of 40 is one SD below the mean. The standardized T-score is reported as the final score for each participant for each domain. For the Global Satisfaction with Sex Life domain, a raw summed score of 6 converts to a T-score of 48.15 with a standard error (SE) of 3.52 (see scoring table in Appendix B). Thus, the 95% confidence interval around the observed score ranges from 41.25 to 55.04 (T-score \pm (1.96*SE) or $48.15 \pm (1.96*3.52)$).

Section 2: Creation of a Customized Sexual Function and Satisfaction Assessment

Introduction

Section 1 was intended to help researchers who are content to use a brief “off the shelf” profile measure of sexual function and satisfaction—the PSxFBP. Section 2 provides information necessary for those users who wish to select specific subdomains and/or specific items within subdomains to create a customized assessment of sexual function and satisfaction using the PROMIS system. Section 2 also provides more detailed information concerning the development, reliability, and validity of the PROMIS SexFS.

Instrument Descriptions

Through the PROMIS Cancer Supplement, instruments assessing multiple components of sexual functioning were developed. Together, these instruments are known as the PROMIS Sexual Function and satisfaction measure (PROMIS SexFS). Some instruments are gender specific. Most items are not specific to cancer, but have thus far only been validated in cancer populations. (Research is ongoing to expand development of the PROMIS SexFS instruments beyond cancer.) The PROMIS SexFS uses a 30-day recall period. Where possible, items use response options common to other PROMIS banks. Some PROMIS SexFS instruments include items from other sexual function instruments, such as the Female Sexual Function Index and the UCLA Prostate Cancer Index.

Available Instruments

PROMIS has 11 sexual function and satisfaction instruments. Five of these instruments are calibrated item banks (e.g. PROMIS Bank v1.0 - Global Satisfaction w Sex Life). This means that if one or more items from within that instrument are administered, a respondent’s score will be calculated using item response theory statistics. If these instruments are administered outside of Assessment Center you may rely on raw score/scale score look-up tables to determine scores (see tables in Appendix C).



Six of the instruments do not have calibrated items (e.g. PROMIS Pool v1.0 - Sexual Activities). This means that items within those instruments are not combined in any way to create a score. Each item in these instruments measures a very specific construct corresponding only to that item (e.g., how much radiation burns have affected one’s satisfaction with their sex life). For any given item in these uncalibrated instruments, the researcher can use the raw item responses directly for analyses. The available instruments are listed in Table 1.

Subdomain Definitions

Definitions for those subdomains measured by the PSxFBP (Global Satisfaction with Sex Life, Interest in Sexual Activity, Lubrication, Vaginal Discomfort, Erectile Function, and Orgasm) are found in Section 1. Definitions for the remaining 5 subdomains are below.

Interfering Factors is a collection of items each of which assesses the person’s perception of the degree to which various factors affect satisfaction with sex life. These factors include symptoms of disease

and side effects from treatment and other issues that have been identified by patients. These items are intended to be “stand alone” items and do not comprise a unidimensional scale. Some items are gender-specific.

Therapeutic Aids is a collection of items each of which assesses the use of hormones, personal lubrications, medications, or devices intended to allow for or improve sexual function. These items are intended to be “stand alone” items and do not comprise a unidimensional scale.

Sexual Activities is a collection of items each of which assesses the frequency of engaging in specific intimate or sexual behaviors either alone or with a partner. These items are intended to be “stand alone” items and do not comprise a unidimensional scale. Some items are gender-specific.

Anal Discomfort is an evaluation of anal irritation, pain, or bleeding during or after anal sex. Items are only asked of people who indicate in the activities subdomain they have had anal sex in the past 30 days. There have not been enough data collected to do psychometric evaluation of these items.

Sexual Function Screener Items ask about sex (gender), whether people are in a relationship that could involve sexual activity, and whether they have had any type of sexual activity with a partner in the past 30 days.

In addition, there are male-specific items related to the Orgasm subdomain that ask about timing of ejaculation and pain or burning during or after ejaculation. These can be administered and scored as single items.

Instrument Name	# of items	Calibrated or Uncalibrated
Global Satisfaction with Sex Life	7	Calibrated
Interest in Sexual Activity	4	Calibrated
Lubrication	8	Calibrated
Vaginal Discomfort	10	Calibrated
Erectile Function	8	Calibrated
Orgasm	3	Uncalibrated
Interfering Factors	10	Uncalibrated
Therapeutic Aids	9	Uncalibrated
Sexual Activities	12	Uncalibrated
Anal Discomfort	5	Uncalibrated
Sexual Function Screener Items	3	Uncalibrated

Table 1: PROMIS Sexual Function and Satisfaction Instruments



Selecting Appropriate Items

All items in the PROMIS SexFS were not intended to be administered together. Researchers should select the sexual function and satisfaction items that are relevant to their specific sample. Some examples are provided.

Example 1: A study proposes to compare three treatment approaches for early stage cervical cancer: surgery alone, surgery and radiation, and radiation alone. In addition to disease control, cancer treatment comorbidities are being compared, including sexual function outcomes. The researchers want to measure key domains of function, including overall sexual satisfaction, interest, vaginal irritation or pain, orgasm, and lubrication. They are also interested in which side effects from treatments affect participants' sex lives, as each of the treatment modalities carries different potential changes in sexual function; surgery usually results in a foreshortened vaginal canal and radiation may cause vaginal mucosal thinning, vaginal adhesions, decreased lubrication and vaginal stenosis. The 10-item **PSxFBP** for women can be used to assess sexual function broadly and distinguish between sexual side effects associated with treatment modality, and can be used to help patients make informed treatment decisions. Additional items on surgical scars, pain, and fatigue from the **Interfering Factors** instrument can help the researchers determine which side effects affect satisfaction with sex life for their participants. Finally, the researchers include the items for women that assess use of **Therapeutic Aids** to determine whether using personal lubricants or hormones modifies sexual satisfaction or function.

Example 2: A study designed to promote compliance with SSRI antidepressants proposes to assess whether sexual function contributes to non-compliance. Patients prescribed fluoxetine are longitudinally followed with monthly assessments of sexual function and frequency of sexual activities in order to determine the relationship between sexual dysfunction and non-compliance. The researchers have room for about 20 items on sexual function, so they choose to use the **PSxFBP** for men (8-items) and women (10-items) to gauge function plus the 12 items from the **Sexual Activities** subdomain. Thus, for all participants in the study, sexual activities, interest in sexual activity, orgasm, and global satisfaction with sex life are assessed. For women, lubrication and vaginal discomfort are also assessed, and for men, erectile function is also assessed.

Example 3: A study of soy-derived estrogen is tested to determine if it improves sexual function among menopausal women self-identified as having hyposexual desire. The researchers choose to administer all items from the **Interest in Sexual Activity** instrument, since sexual desire is their main outcome of interest. They also administer the **PSxFBP** for women to assess satisfaction with sex life, lubrication, vaginal discomfort, and orgasm.

A Note on Response Options for Sexual Activities

Most sexual activity items are available using two different sets of responses. Items identified with an "a" in their Item ID use the response options 1=Have not done in the past 30 days, 2=Once, 3=Two to three times, 4=Four to five times, 5=Six or more times. Items identified with a "b" in their Item ID use the response options 1=Have not done in the past 30 days, 2=Once a week or less, 3=Once every few days, 4=Once a day, 5=More than once a day. As you can see, the "a" response options reflect less activity. This set of response options is likely most appropriate for individuals for whom you expect reduced sexual activity (e.g., cancer patients receiving chemotherapy). The "b" response options reflect higher levels of sexual activity. This set of response options would be most appropriate for individuals you expect to have higher levels of sexual activity (e.g., healthy individuals). Investigators should carefully consider their purpose in recording sexual activities and select response options that are most appropriate. It is possible that the "a" and "b" options available here are not the best for a particular research setting. Investigators might also consider whether a daily sexual activity log could be used in place of these items, which require a 30-day recall period.



Procedures and Data in Support of Validity and Reliability

Face Validity. Face validity is established when subject matter experts agree that the scale appears to measure its intended focus. Face validity for the PROMIS SexFS scales was established with a review by expert panels within and external to the PROMIS SexFS committee; all experts concurred that the items within the scales appeared to measure sex function.

Content Validity. Content validity refers to how well the scale assesses all aspects of the construct being measured. Establishing the content validity of PROMIS instruments began with patient input to assure that the subdomains and their items corresponded to reported patient experiences, and with a review by expert panels to assure that the selected theoretical constructs corresponded to the scientific literature. Using a consensus-driven approach, the PROMIS SexFS committee conducted a literature search for articles published from 1991 through 2007, yielding 257 articles that reported the administration of a psychometrically evaluated sexual function measure to individuals diagnosed with cancer. With few exceptions, the 31 identified measures had not been widely tested in cancer populations (Jeffery et al., 2009). We collated available items from the measures and created preliminary domain definitions. Each item was then subjected to detailed review to eliminate repetition within bins (“winnowing”) and to develop uniform recall periods and response categories. After qualitative expert item review, 47 extant items were selected for further testing. Concurrently, we conducted 16 focus groups with 109 cancer patients (Flynn et al., 2010). These groups explored the impact of cancer and its treatments on sexual experience to determine whether domain definitions and the identified items reflected patients’ personal experiences. Separate focus groups were held for patients in active treatment for breast, prostate, lung, colorectal, gynecological, and other (mixed) cancers and for survivors after treatment for breast, prostate, gynecological, and other cancer types. We developed a matrix of themes and groups, which was double-coded (inter-rater reliability was over 90%). As a check on the data we received from the patient focus groups, we conducted 2 clinician focus groups to assess the clinical relevance of the proposed conceptual model and obtain clinicians’ views of how cancer and its treatment affected patients’ sexual health. New items were created to address conceptual gaps identified by the focus group participants. With updated items in hand, we conducted cognitive interviews with patients (n=39) (Fortune-Greeley et al., 2009). Each item was seen by at least 5 patients, at least 1 of whom was not white and at least 2 of whom had less than a 9th grade reading level. 87 items were passed through to the next phase. We convened 7 experts on sexual function and cancer to review this work to date.

The item-testing phase consisted of large-scale data collection (n=819; 388 males, 430 females, 1 person did not specify sex) and administration of the items in national and local samples through the NexCura Internet Panel, the Duke University tumor registry, and the Duke oncology clinics.

(Appendix D shows patient characteristics, including the distribution of cancer types.) We also added targeted recruitment of additional lesbian, gay, and bisexual cancer patients and survivors through online communities. Psychometric

analysis of the items followed established PROMIS methodology (Reeve et al., 2007) and resulted in 11 instruments: 5 calibrated and 6 uncalibrated. A summary of fit statistics are shown in Table 2.

Instrument Name	CFI	TLI	RMSEA
Global Satisfaction with Sex Life	0.983	0.976	0.168
Interest in Sexual Activity	0.998	0.995	0.129
Lubrication	0.985	0.979	0.187
Vaginal Discomfort	0.993	0.991	0.124
Erectile Function	0.988	0.986	0.134
CFI: comparative fit index			
TLI: Tucker-Lewis index			
RMSEA: root-mean-square error of approximation			

Table 3: Fit Indices for Confirmatory Factor Analysis of Calibrated Subdomains.

Women (N = 430)		Men (N = 388)	
Measures	r*	Measures	r*
PROMIS Interest in Sexual Activity	0.84	PROMIS Interest in Sexual Activity	0.82
FSFI [†] Desire	(0.82)	IIEF [‡] Desire	(0.79)
FSFI Arousal	0.71		
	(0.68)		
PROMIS Lubrication	0.92	PROMIS Erectile Function	0.81
FSFI Lubrication	(0.9)	IIEF Erectile Function	(0.69)
PROMIS Vaginal Discomfort	0.9		
FSFI Pain	(0.84)		
PROMIS Orgasm	0.78	PROMIS Orgasm	0.62
FSFI Orgasm	(0.78)	IIEF Orgasmic Function	(0.62)
PROMIS Global Satisfaction with Sex Life	0.76	PROMIS Global Satisfaction with Sex Life	0.82
FSFI Satisfaction	(0.62)	IIEF Overall Satisfaction	(0.66)
		IIEF Intercourse Satisfaction	0.75
			(0.68)

Note: Correlations in parentheses are for the PROMIS Sexual Function Brief Profile scores.
^{*}Pearson correlation coefficients
[†]Female Sexual Function Index
[‡]International Index of Erectile Function

Table 2: Correlations between PROMIS Sexual Function and Satisfaction Subdomains and Corresponding Measures.

Construct Validity. Construct validity refers to how well scores on the measure are related to other variables that, for theoretical reasons, ought to be related to the measure in question. Construct validity of the PROMIS SexFS has been assessed in two ways.

First, we used data from the 819 patients with cancer (see above) to examine the correlations between subdomains of the PROMIS SexFS and other measures of similar constructs. These are displayed in Table 3. In general, these correlations provide strong evidence for the construct validity of the PROMIS SexFS.

Second, we examined whether scores on selected subdomains of the PROMIS SexFS could discriminate between groups that should, in theory, differ in terms of their sexual experiences. During item testing, participants were also asked whether they had ever asked an oncology professional about sexual problems. We hypothesized that asking for help with sexual problems may indicate a clinically meaningful decrement in function. As Table 4 shows, those who had asked for help had significantly greater interest in sexual activity and increased vaginal discomfort and lower levels of erectile function, lubrication, orgasm, and overall satisfaction. Furthermore, the differences were as high as three-quarters of a standard deviation. These effect sizes were greater than or equal to the effects for the corresponding subscales of the FSFI and IIEF. In three cases, the PROMIS SexFS and PSxFBP detected statistically significant ($p < .05$) differences between those who did and did not ask, whereas the FSFI or IIEF did not.

Reliability. Two types of reliability data are available at this time for the PROMIS SexFS. First, estimates of internal consistency (Cronbach’s alpha) were computed for all calibrated banks. They are displayed in Table 4.

PROMIS Sexual Function and Satisfaction Measure			PROMIS SexFS Brief Profile		Legacy Measures		
Domain	Effect Size	p-value ^b	Effect Size	p-value ^b	Domain	Effect Size	p-value ^b
Interest in Sexual Activity (women)	0.18	0.16	0.16	0.19	FSFI Desire	-0.05	0.69
					FSFI Arousal	-0.13	0.35
Interest in Sexual Activity (men)	0.22	0.04	0.21	0.05	IIEF Desire	0.14	0.2
Erectile Function	-0.45	<.0001	-0.49	<.0001	IIEF Erectile Function	-0.22	0.04
Lubrication	-0.75	<.0001	-0.66	<.0001	FSFI Lubrication	-0.67	<.0001
Vaginal Discomfort	0.75	<.0001	0.6	<.0001	FSFI Pain	-0.6	<.0001
Orgasm (women)	-0.4	<.0001	-0.4	0.003	FSFI Orgasm	-0.11	0.41
Orgasm (men)	-0.55	<.0001	-0.55	<.0001	IIEF Orgasmic Function	-0.55	<.0001
Global Satisfaction with Sex Life (women)	-0.24	0.05	-0.27	0.03	FSFI Satisfaction	0.08	0.54
Global Satisfaction with Sex Life (men)	-0.15	0.15	-0.18	0.09	IIEF Overall Satisfaction	-0.16	0.14

Abbreviations: PROMIS Patient Reported Outcomes Measurement Information System; FSFI Female Sexual Function Index; IIEF International Index of Erectile Function
a: Difference in the means between people who answered “Yes” (N = 237) and “No” (N = 569) to the question, “Have you ever asked an oncology doctor or nurse about problems with your sex life?” divided by the common standard deviation.
b: From t-test comparing PROMIS measure to legacy measure.

Table 4: Effect Sizes Discriminating Askers From Non-Askers^a (N=806).



All indicate excellent internal consistency. Second, test-retest reliability was examined in a sample of 202 participants (101 male, 101 female), about half of whom had some chronic disease. Participants completed the PROMIS SexFS twice with one month between test administrations. Intraclass correlation coefficients between the two administrations are shown in Table 5, ranging from .71 - .87.

Measures	Women		Men	
	Cronbach's alpha (N = 430)	ICC PROMIS SxeFS (PSxFBP) (N = 101)	Cronbach's alpha (n = 388)	ICC PROMIS SxeFS (PSxFBP) (N = 101)
PROMIS Interest in Sexual Activity	0.89	0.77 (0.72)	0.87	0.71 (0.65)*
PROMIS Lubrication	0.95	0.87 (0.87)		
PROMIS Vaginal Discomfort	0.94	0.80 (0.75)		
PROMIS Erectile Function			0.92	0.87 (0.77)
PROMIS Global Satisfaction with Sex Life	0.93	0.75 (0.69)	0.92	0.74 (0.66)

*Numbers reflect the deletion of a single outlier with a "5" at first administration and a "1" at second administration. Inclusion of the outlier results in ICCs of 0.55 and 0.54 for the full bank and PSxFBP, respectively.

Table 5: Reliability of Calibrated Subdomains.

Procedures for Selecting the PROMIS Sexual Function and Satisfaction Profile in Assessment Center

The PROMIS Sexual Function and Satisfaction Profile is a publicly available instrument in the Assessment Center library. Assessment Center allows you to create a study-specific data collection website for capturing participant data. There are three versions of the Profile within Assessment Center: a male version, a female version and a combined gender version which is appropriate for males and females. The combined gender version will branch the respondent to the appropriate questions (erectile function versus lubrication and vaginal discomfort) based on gender.

Before using an existing PROMIS Sexual Function and Satisfaction instrument, or creating a custom one, you must first create a study. To do this, select the Studies tab and click on the Create New Study button. Enter study information and click on the Save button. Select the Studies tab and the new study should appear in the My Studies box. To add a PROMIS Sexual Function and Satisfaction Profile to your study in Assessment Center first, navigate to the Study Content page by clicking the Instruments tab. Next, click the Add button to access the Add an Instrument page. From this page, you can search for sexual function and satisfaction instruments by using the search drop lists at the top of the page. Once you have identified the appropriate PROMIS Sexual Function and Satisfaction Profile instrument, check the box next to the desired instrument, and click Add to Study button at the top or bottom of the search results box. Additional information about using Assessment

Center is available in the Assessment Center User Manual (available at assessmentcenter.net) or within the application through Help (upper right corner hyperlink).

Procedures for Creating a Custom PROMIS Sexual Function and Satisfaction Instrument in Assessment Center

Assessment Center allows you to create a custom instrument so you may individually select sexual function and satisfaction items of interest to administer to participants. To do this, navigate to the Study Content page by clicking the Instruments tab. Click on the Create button. Enter instrument information on the Instrument Properties page selecting Short Form for Instrument Type. Click Save to be navigated back to the Study Content page. Next, click on your newly created instrument's name which will appear as a hyperlink to be navigated to the Instrument Details page. Then click on the Find Items button to access the Add an Item page. From this page, you can search for sexual function and satisfaction instruments by using the search drop lists at the top of the page. Once you have identified the appropriate PROMIS Sexual Function and satisfaction instrument (e.g., Therapeutic Aids), click on the plus sign to the left of the instrument name. The page will expand to display all items within the instrument. Check the box next to the desired items and click Add to Instrument at the top or bottom of the search results box. To view your custom instrument with the items you have just selected, click on the Instruments tab. Next click the plus sign to the left of the instrument name.

Procedures for Previewing PROMIS Studies in Assessment Center

Before launching a study, Assessment Center allows you to first preview the study. To preview your study you must first click on the Team hyperlink to the right of the desired study. Next identify the study team members. Assign roles to individual members by highlighting their name and checking the box next to the desired role (*Note: the team member launching the study preview must have the role of Study Administrator or Instrument/Item Administrator*). Next select the Preview tab. Click on the Preview Study button (*Note: the preview may take a few seconds to launch*). Next click on the Continue button and follow instructions on the study Welcome page.



The following experts generously provided helpful input to the PROMIS sexual function and satisfaction domain working group:

	<u>Sexual Function and Satisfaction Domain Group (including NIH)</u>	<u>Duke Clinical Research Institute CCGE staff</u>	<u>Duke University School of Nursing</u>
Erick Janssen, PhD, MA Ray Rosen, PhD Stacy Tessler Lindau, MD, MAPP Jeanne Carter, PhD Michael Perelman, PhD Leslie R. Schover, PhD John P. Mulhall, MD David M. Latini, PhD Barbara L. Andersen, PhD Sara I. McClelland, PhD	Amy Abernethy, MD Joan Broderick, PhD Deborah Bruner, PhD, RN Jill Cyranowski, PhD Susan Czajkowski, PhD Elizabeth Hahn, MA Diana Jeffery, PhD Francis Keefe, PhD Jin-Shei Lai, PhD Richard Luecht, PhD Susan Magasi, PhD Laura Porter, PhD Jennifer Reese, PhD Bryce Reeve, PhD Rebecca Shelby, PhD Ashley Wilder, PhD, MPH	Carrie Dombeck, MA Maria Fawzy Alice Fortune-Greeley Angel Moore, MSPH Damon Seils Janice Tzeng	Lucy Andrzejewski Teresa Baker Henry Beresford Monie Clayton Teresa Ebel Linda Folsom Mindy Kash Patrick Lane Diane Langley Justin Levens Denise Snyder, MS, RD, CSO, LDN Valeda Stull Megan Williams, MSW, MSPH

Contact Us

For more information about PROMIS, accessing the PROMIS Sexual Function instruments or administering them through Assessment Center, contact us at help@healthmeasures.net.

References

DeWalt, D., Rothrock, N., Yount, S., Stone, A. A., & on behalf of the PROMIS cooperative group. (2007). Evaluation of item candidates: the PROMIS qualitative item review. *Medical Care, 45(5)*, S12-21.

Fortune-Greeley, A. K., Flynn, K. E., Jeffery, D. D., Williams, M. S., Keefe, F. J., Reeve, B. B., Willis, G. B., & Weinfurt, K. P. (2009). Using cognitive interviews to evaluate items for measuring sexual functioning across cancer populations: improvements and remaining challenges. *Quality of Life Research, 18(8)*, 1085-93.

Jeffery, D. D., Tzeng, J. P., Keefe, F.J., Porter, L. S., Hahn, E. A., Flynn, K. E., Reeve, B. B., & Weinfurt, K. P. (2009). Initial report of the cancer Patient-Reported Outcomes Measurement Information System (PROMIS) sexual function committee: Review of sexual function measures and domains used in oncology. *Cancer, 115(6)*, 1142-53.

Cella D, Riley W, Stone A, Rothrock N, Reeve B, Yount S, Amtmann D, Bode R, Buysse D, Choi S, Cook K, DeVellis R, DeWalt D, Fries JF, Gershon R, Hahn EA, Lai JS, Pilkonis P, Revicki D, Rose M, Weinfurt K, Hays R. (2010). The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005-2008. *J Clin Epidemiol. 63(11)*, 1179-84.



Flynn, K. E., Jeffery, D. D., Keefe, F. J., Porter, L. S., Shelby, R. A., Fawzy, M. R., Gosselin, T. K., Reeve, B. B., & Weinfurt, K. P. (2011). Sexual functioning along the cancer continuum: focus group results from the Patient-Reported Outcomes Measurement Information System (PROMIS). *Psycho-Oncology, 20(4)*, 378-86.

Flynn K.E., Reese J.B., Jeffery D.D., Abernethy A.P., Lin L., Shelby R.A., Porter L.S., Dombeck C.B., Weinfurt K.P. (in press). Patient experiences with communication about sex during and after treatment for cancer. *Psycho-Oncology*.

Flynn K.E., Jeffery D.D., Reeve B.B., Lin L., Wilder Smith A., Abernethy A.P., Reese J.B., Weinfurt K.P. "Progress on the PROMIS® Sexual Function Measure." Poster presented at the International Society for Quality of Life Research Annual Conference; October 29, 2010; London, England.

Weinfurt, K. "Improving the Measurement of Sexual Outcomes in Cancer: The NIH PROMIS Approach." Center for Outcomes and Policy Research, Dana-Farber Cancer Institute, Boston, Massachusetts, May 17, 2011

Flynn K.E., Weinfurt K.P. "Development and Validation of the PROMIS Sexual Function Measure." Presentation at the Duke Translational Medicine Institute Annual Research Career Day; May 20, 2011; Durham, NC, USA.

Flynn K.E. "Assessing Sexual Health in Cancer Survivors." Presentation at the Cancer Survivorship and Sexual Health Symposium sponsored by the International Society for Sexual Medicine and the Sexual Medicine Society of North America; Washington, DC, USA; June 18, 2011.

Weinfurt K.P., Lin L., Broderick J.E., Dombeck C.B., Fawzy, M.R., Snyder D.C., Williams M.S., Flynn K.E. "Validity of 1-month Recall for Components of Sexual Function." To be presented as a poster at the International Society for Quality of Life Research Annual Conference; October 28, 2011; Denver, Colorado, USA.

Flynn K.E., Lin L., Dombeck C., Fawzy M., Abernethy A.P., Bruner D.W., Reese J.B., Reeve B.B., Smith A.W., Weinfurt K.P. "Validating the PROMIS® Sexual Function Brief Profile Measures." To be presented at the International Society for Quality of Life Research Annual Conference; October 29, 2011; Denver, Colorado, USA.

Weinfurt K.P., Lin L., Broderick J.E., Dombeck C.B., Fawzy, M.R., Snyder D.C., Williams M.S., Flynn K.E. "Mood and Gender Effects on the Accuracy of 30-day Recall of Sexual Function." To be presented at the International Society for Quality of Life Research Annual Conference; October 29, 2011; Denver, Colorado, USA.



Appendix B - Sexual Function and Satisfaction Measures Brief Profile Look-up Tables

Global Satisfaction with Sex Life <i>Score Conversion Table</i>		
Raw Score	T Score	SE*
2	30.67	4.86
3	36.8	3.84
4	40.94	3.59
5	44.76	3.6
6	48.15	3.52
7	51.5	3.57
8	55.11	3.54
9	58.98	3.76
10	65.6	5.23

*SE=Standard Error on T-score metric

Interest in Sexual Activity <i>Score Conversion Table</i>		
Raw Score	T Score	SE*
2	33.42	4.72
3	40.01	2.82
4	43.64	2.69
5	47.46	2.82
6	51.16	2.8
7	54.86	2.85
8	58.96	2.87
9	63.28	3.06
10	69.97	4.37

*SE=Standard Error on T-score metric

Lubrication <i>Score Conversion Table</i>		
Raw Score	T Score	SE*
2	37.05	5.45
3	43.58	3.26
4	46.25	3.07
5	48.5	2.99
6	50.64	2.96
7	52.84	2.98
8	55.3	3.06
9	58.55	3.38
10	64.82	5.24

*SE=Standard Error on T-score metric

Vaginal Discomfort <i>Score Conversion Table</i>		
Raw Score	T Score	SE*
3	34.34	5.3
4	41.13	3.56
5	45.4	2.83
6	48.09	2.69
7	50.51	2.61
8	52.62	2.61
9	54.55	2.63
10	56.5	2.69
11	58.56	2.78
12	61.03	2.99
13	64.32	3.42
14	69.81	4.96

*SE=Standard Error on T-score metric

Erectile Function <i>Score Conversion Table</i>		
Raw Score	T Score	SE*
3	36.84	5.41
4	42.81	3.22
5	44.88	3.05
6	46.76	2.82
7	48.44	2.65
8	49.99	2.58
9	51.51	2.58
10	53.08	2.6
11	54.78	2.64
12	56.64	2.66
13	58.69	2.72
14	61.32	2.96
15	67.25	4.67

*SE=Standard Error on T-score metric



Appendix C - Sexual Function and Satisfaction Bank Look-up Tables

Vaginal Discomfort		
Score Conversion Table		
Raw Score	T Score	SE*
10	33.20	5.05
11	38.71	3.26
12	41.09	2.98
13	42.68	2.67
14	44.01	2.43
15	45.13	2.19
16	46.11	2.01
17	46.96	1.91
18	47.74	1.86
19	48.47	1.83
20	49.16	1.80
21	49.83	1.78
22	50.48	1.77
23	51.12	1.77
24	51.74	1.77
25	52.35	1.77
26	52.96	1.77
27	53.56	1.79
28	54.17	1.79
29	54.77	1.80
30	55.38	1.81
31	56.00	1.84
32	56.63	1.86
33	57.28	1.89
34	57.95	1.93
35	58.66	2.00
36	59.40	2.08
37	60.20	2.17
38	61.04	2.28
39	61.93	2.41
40	62.87	2.55
41	63.88	2.70
42	64.96	2.88
43	66.14	3.09
44	67.42	3.29
45	68.89	3.57
46	70.34	3.74
47	72.36	4.09
48	74.03	4.31
49	77.08	4.90

*SE= Standard Error on T-score

Lubrication		
Score Conversion Table		
Raw Score	T Score	SE*
8	30.99	4.74
9	35.47	3.16
10	37.54	2.72
11	38.97	2.48
12	40.14	2.27
13	41.16	2.10
14	42.06	1.98
15	42.89	1.90
16	43.66	1.85
17	44.40	1.80
18	45.09	1.78
19	45.78	1.77
20	46.45	1.77
21	47.12	1.76
22	47.78	1.77
23	48.46	1.79
24	49.14	1.80
25	49.84	1.81
26	50.54	1.82
27	51.26	1.84
28	51.99	1.86
29	52.74	1.87
30	53.52	1.88
31	54.32	1.90
32	55.16	1.92
33	56.05	1.96
34	57.01	2.01
35	58.06	2.10
36	59.26	2.25
37	60.67	2.50
38	62.38	2.82
39	64.78	3.33
40	69.26	4.85

*SE= Standard Error on T-score

Erectile Function		
Score Conversion Table		
Raw Score	T Score	SE*
8	30.72	5.42
9	33.94	5.38
10	32.59	5.62
11	34.48	5.35
12	34.19	5.18
13	38.05	3.83
14	39.75	3.72
15	40.68	3.90
16	42.28	3.16
17	43.57	2.86
18	44.65	2.69
19	45.66	2.55
20	46.59	2.45
21	47.46	2.38
22	48.30	2.33
23	49.11	2.30
24	49.89	2.29
25	50.68	2.29
26	51.46	2.30
27	52.26	2.32
28	53.08	2.37
29	53.93	2.42
30	54.83	2.50
31	55.79	2.62
32	56.80	2.71
33	57.94	2.92
34	59.27	3.29
35	61.08	4.00
36	62.12	3.75
37	64.00	3.67
38	68.47	4.76

*SE= Standard Error on T-score

Global Satisfaction with Sex Life		
Score Conversion Table		
Raw Score	T Score	SE*
7	29.59	4.56
8	34.45	3.10
9	37.14	2.73
10	39.16	2.54
11	40.77	2.42
12	42.15	2.33
13	43.40	2.26
14	44.55	2.21
15	45.63	2.18
16	46.66	2.16
17	47.66	2.16
18	48.63	2.16
19	49.60	2.16
20	50.56	2.17
21	51.53	2.18
22	52.51	2.18
23	53.49	2.18
24	54.49	2.18
25	55.49	2.17
26	56.51	2.17
27	57.54	2.18
28	58.61	2.20
29	59.73	2.23
30	60.93	2.31
31	62.25	2.43
32	63.74	2.60
33	65.54	2.90
34	67.85	3.33
35	72.01	4.63

*SE= Standard Error on T-score

Interest in Sexual Activity		
Score Conversion Table		
Raw Score	T Score	SE*
4	32.03	4.78
5	37.01	3.41
6	40.15	2.94
7	42.55	2.73
8	44.67	2.74
9	46.79	2.80
10	48.87	2.81
11	50.86	2.78
12	52.85	2.79
13	55.15	2.84
14	57.58	2.81
15	59.78	2.83
16	62.19	2.94
17	64.98	3.07
18	68.43	3.36
19	71.76	3.92
20	76.17	4.87

*SE= Standard Error on T-score

Appendix D. Characteristics of Validation Sample

Characteristic	Total (N = 819)
Age, mean ± SD, y	58.5 ± 11.8
Age group, No. (%)	
≤ 40 years	59 (7)
41 to 50 years	127 (16)
51 to 64 years	377 (46)
65 to 79 years	232 (28)
≥ 80 years	21 (3)
Race, No. (%)	
Black or African American	80 (10)
American Indian/Alaska Native	10 (1)
Asian	12 (1)
Native Hawaiian/Other Pacific Islander	10 (1)
White	705 (87)
Multiple races or other	2 (< 1)
Hispanic or Latino ethnicity, No. (%)	21 (3)
Educational attainment, No. (%)	
Less than high school	21 (3)
High school graduate/GED	100 (12)
Some college	255(31)
College degree	229 (28)
Advanced degree (MA, PhD, MD)	211 (26)
Treatment status in past month, No. (%)	
None (ie, posttreatment follow-up)	526 (64)
Undergoing treatment	290 (36)
Radiation therapy	29 (10)
Hormonal therapy (eg, tamoxifen, anastrozole, leuprolide)	140 (48)
Chemotherapy (injection or oral)	116 (40)
Immunotherapy (eg, interferon)	9 (3)
Other	36 (12)
Recurrence of cancer, No. (%)	151 (18)
Cancer spread to lymph nodes, No. (%)	202 (25)
Cancer spread to another area, No. (%)	134 (16)
Primary cancer diagnosis, No. (%)	
Bone/muscle cancer	14 (2)
Brain cancer	4 (< 1)
Breast cancer	252 (35)
Colorectal	98(13)
Esophageal or stomach cancer	17 (2)
Gynecologic cancer	29 (4)
Head/neck cancer	9 (< 1)
Hodgkin lymphoma	23 (3)
Leukemia	20 (3)
Liver cancer	3 (< 1)
Lung cancer	56 (8)
Melanoma	4 (< 1)
Multiple Myeloma	2 (< 1)
Non-Hodgkin lymphoma	12 (2)
Pancreatic cancer	5 (< 1)
Prostate cancer	146 (20)
Urologic cancer	23 (3)