

PAIN QUALITY MEASURE DIFFERENCES

A brief guide to differences between the PROMIS® Pain Quality instruments:

ADULT	PEDIATRIC
PROMIS Scale v2.0 – Neuropathic Pain Quality 5a PROMIS Scale v2.0 – Nociceptive Pain Quality 5a PROMIS Pool v1.0 – Pain Quality*	PROMIS Pediatric Bank GenPop v3.0 – Pain Quality PROMIS Pediatric Bank GenPop v3.0 – Pain Quality (screen-to-CAT) PROMIS Pediatric Short Form GenPop v3.0 – Pain Quality – Affective 8a PROMIS Pediatric Short Form GenPop v3.0 – Pain Quality – Sensory 8a PROMIS Pediatric Bank v2.0 – Pain Quality* PROMIS Pediatric Short Form v2.0 – Pain Quality – Affective 8a* PROMIS Pediatric Short Form v2.0 – Pain Quality – Sensory 8a*

*Retired measure

ABOUT PAIN QUALITY

The adult PROMIS Pain Quality scales assess self-reported nociceptive pain and neuropathic pain.

- Nociceptive pain is caused by stimulation of peripheral nerve fibers (nociceptors) in the context of a normally functioning somatosensory nervous system. The PROMIS Nociceptive Pain Scale includes descriptors of pain sensations, such as “achy”, “deep”, “sore”, and “tender”, commonly reported by people with some qualities typical of nociceptive pain.
- Neuropathic pain is caused by damage to the peripheral somatosensory nervous system, part of the nervous system involved in bodily feelings. This damage can be caused by an abnormality, trauma or disease. The PROMIS Neuropathic Pain Scale asks to what degree (from “not at all” to “very much”) the respondent’s pain shows qualities typical of neuropathic pain (i.e., felt “numb”, “tingly”, like “pins and needles”, “stinging”, or “electrical”).

The pediatric Pain Quality measures assess specific physical sensations and affective components associated with pain. Because pain can be felt and described in so many ways, this category of pain contains a variety of attributes, such as perceived temperature (e.g., cold), sensations (e.g., throbbing), and perceived affective qualities of pain (e.g., uncomfortable).

The Pain Quality measures are universal rather than disease-specific. All assess pain quality over the past seven days. Pain quality instruments are available for adults (ages 18+) and pediatric self-report (ages 8-17).

INTRODUCTION TO ASSESSMENT OPTIONS

There is one administration option for assessing adult Pain Quality: fixed length scales. No computer adaptive test (CAT) is available. When administering a scale, instruct respondents to answer all of the items (i.e., questions or statements) presented.



There are two administration options for assessing pediatric Pain Quality: short forms and computer adaptive tests (CATs). When administering a short form, instruct participants to answer all of the items (i.e., questions or statements) presented. With a CAT, participant responses guide the system's choice of subsequent items from the full item bank (56 items in total). Although items differ across respondents taking a CAT, scores are comparable across participants.

Some administrators may prefer to ask the same question of all respondents or of the same respondent over time, to enable a more direct comparability across people or time. In these cases, or when paper administration is preferred, a short form would be more desirable than a CAT. This guide provides information on all pediatric Pain Quality short forms and a CAT instrument.

CAT: A minimum number of items (e.g., 4) must be answered in order to receive a score for Pain Quality CATs. The response to the first item will guide the system's choice of the next item for the participant. The participant's response to the second item will dictate the selection of the following question, and so on. As additional items are administered, the potential for error is reduced and confidence in the respondent's score increases. CATs will continue until either the standard error drops below a specified level (e.g., on the T-score metric 3.0), or the participant has answered the maximum number of questions (e.g., 12), whichever occurs first. For some CATs, specifically "recommended" and "screen-to-CAT" there are additional stopping rules. These include stopping when the standard error isn't improving much or if a respondent is asymptomatic. For details on the exact stopping rules for Pain Quality CATs, see below.

CAT versus Short Form: Whether one uses a short form or CAT, the score metric is Item Response Theory (IRT), a family of statistical models that link individual questions to a presumed underlying trait or concept of pain quality represented by all items in the item bank. When choosing between a CAT and a short form, it is useful to consider the demands of computer-based assessment, and the psychological, physical, and cognitive burden placed on respondents as a result of the number of questions asked.

VERSION DIFFERENCES

Some PROMIS domains have multiple versions of instruments (i.e., v1.0, v1.1, v2.0, v3.0). Generally, **it is recommended that you use the most recent version available which can be identified as the instrument with the highest version number**. In most cases, an instrument that has a decimal increase (v1.0 to v1.1) retains the same item-level parameters as well as instrument reliability and validity. In cases where a version number increases by a whole number (e.g., v1.0 to v2.0), the changes to the instrument are more substantial. PROMIS initially produced Pain Quality item pool v1.0 for adults. This was a collection of items that were not calibrated but instead intended to be used descriptively or for further research. The v2.0 Neuropathic and Nociceptive Pain measures are calibrated and the preferred assessment tool.

Pediatric

Changes from v2.0 to GenPop v3.0

- The GenPop v3.0 Pediatric measures replaced the v2.0 measures.
- The GenPop v3.0 measures were re-normed. This means that the scores produced by v3.0 measures are NOT equivalent to scores from older measures (i.e., v1.0, v1.1, v2.0).
- The v2.0 measures were developed with U.S. children sampled from a combination of the general population and those with chronic conditions. This means a v2.0 T-score of 50 is based on the mean of a sample comprised of a mix of children from the general population AND children with chronic conditions. The re-normed v3 GenPop measures are now purely based on a sample from the general pediatric population. This makes interpreting a PROMIS score easier as it is referencing just the



general population. The use of “GenPop” (general population) is used to convey the difference in metrics between v3.0 and earlier versions of the measures.

- Earlier versions of the PROMIS Pediatric measures can be converted to the v3.0 GenPop metric. Instructions are included in the Pain Quality Scoring Manual.
- No items were revised between the v2.0 and GenPop v3.0 Pain Quality measures.

Pediatric CAT Stopping Rules:

- Measures named “Bank” are administered by default as computer adaptive tests.
- CATs with the same version number (e.g., v2.0 recommended, v2.0 screen-to-CAT) are based on the same item bank but use different stopping rules.

PROMIS Pediatric Bank GenPop v3.0 – Pain Quality stopping rules:

- Minimum number of items administered = 4
- Stop when one of these occurs:
 - 8 items are administered OR
 - Standard error is below 0.3 on the theta metric (3.0 on the T-score metric) OR
 - Standard error changes by less than 0.01 on the theta metric (0.1 on the T-score metric)

PROMIS Pediatric Bank GenPop v3.0 – Pain Quality (screen-to-CAT) stopping rules:

- If the responses to the first two items are both the “healthiest” responses, then stop.
- If the responses to the first two items are NOT the “healthiest” responses, proceed with the PROMIS Pediatric Bank GenPop v3.0 – Pain Quality CAT stopping rules.

PROMIS Pediatric Bank GenPop v2.0 – Pain Quality stopping rules:

- Minimum number of items administered = 4
- Stop when one of these occurs:
 - 12 items are administered OR
 - Standard error is below 0.3 on the theta metric (3.0 on the T-score metric) OR

SCORES

For most PROMIS instruments, a score of 50 is the average for the United States general population with a standard deviation of 10 because calibration testing was performed on a large sample of the general population. You can read more about the calibration and centering samples on HealthMeasures.net (<http://www.healthmeasures.net/score-and-interpret/interpret-scores/promis>). The T-score is provided with an error term (Standard Error or SE). The Standard Error is a statistical measure of variance and represents the “margin of error” for the T-score.

Important: *A higher PROMIS T-score represents more of the concept being measured. For negatively-worded concepts like pain quality, a T-score of 60 is one SD worse than average. By comparison, a pain quality T-score of 40 is one SD better than average.*

STATISTICAL CHARACTERISTICS

There are four key features of the score for Pain Quality:

- **Reliability:** The degree to which a measure is free of error. It can be estimated by the internal consistency of the responses to the measure, or by correlating total scores on the measure from two time points when there has been no true change in what is being measured (for z-scores, reliability = $1 - SE^2$).
- **Precision:** The consistency of the estimated score (reciprocal of error variance).
- **Information:** The precision of an item or multiple items at different levels of the underlying continuum (for z-scores, information = $1/SE^2$).
- **Standard Error (SE):** The possible range of the actual final score based upon the scaled T-score. For example, with a T-score of 52 and a SE of 2, the 95% confidence interval around the actual final score ranges from 48.1 to 55.9 (T-score $\pm (1.96*SE) = 52 \pm 3.9 = 48.1$ to 55.9).

The final score is represented by the T-score, a standardized score with a mean of 50 and a standard deviation (SD) of 10.

PREVIEW OF SAMPLE ITEM

Figure 1 is an excerpt from the paper version of the adult five-item Neuropathic Pain Quality Scale. This is the paper version format used for all Pain Quality instruments.

In the past 7 days...		Not at all	A little bit	Somewhat	Quite a bit	Very much
PAGUAL15r	Did your pain feel sore?	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
PAGUAL05r	Did your pain feel tender?	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

Figure 1

FREQUENTLY ASKED QUESTIONS (FAQs)

Q: I am interested in learning more. Where can I do that?

Review the HealthMeasures website at www.healthmeasures.net.

Q: Are these instruments available in other languages?

No. However, periodically check the HealthMeasures website (<http://www.healthmeasures.net/explore-measurement-systems/promis/intro-to-promis/available-translations/117-available-translations>) for current information on PROMIS translations.

Q: Can I make my own short form?

Yes, custom short forms can be made by selecting any items from an item bank. This can be scored using the Scoring Service (https://www.assessmentcenter.net/ac_scoringervice).