



PROMIS Domain Framework

Respondent	Category	Domain	Sub-Domain	Domain Definition
Physical Health				
Adult (Ages 18+)	Physical Health	Pain – Behavior		Behaviors that typically indicate to others that an individual is experiencing pain. These actions or reactions can be verbal or nonverbal, and involuntary or deliberate.
Adult (Ages 18+)	Physical Health	Pain – Interference		Consequences of pain on relevant aspects of one’s life. This includes the extent to which pain hinders engagement with social, cognitive, emotional, physical, and recreational activities.
Adult (Ages 18+)	Physical Health	Pain Intensity		How much a person hurts
Adult (Ages 18+)	Physical Health	Pain Quality – Neuropathic Pain		Caused by damage to the peripheral somatosensory nervous system, part of the nervous system involved in bodily feelings. This damage can be caused by an abnormality, trauma or disease.
Adult (Ages 18+)	Physical Health	Pain Quality – Nociceptive Pain		Caused by stimulation of peripheral nerve fibers (nociceptors) in the context of a normally functioning somatosensory nervous system.
Adult (Ages 18+)	Physical Health	Fatigue		Range of symptoms, from mild subjective feelings of tiredness to an overwhelming, debilitating, and sustained sense of exhaustion that likely decreases one’s ability to execute daily activities and function normally in family or social roles.
Adult (Ages 18+)	Physical Health	Itch – Interference		General issues related to quality of life impairment from itch (pruritis).
Adult (Ages 18+)	Physical Health	Itch – Quality		Assesses the subjective description of the sensation of itch with a checklist.
Adult (Ages 18+)	Physical Health	Itch – Scratching Behavior		Quality of life impairment from scratching behavior and the physical manifestations of itch (pruritis).
Adult (Ages 18+)	Physical Health	Itch – Severity		Characteristics of itch, including intensity, frequency, and time of occurrence.
Adult (Ages 18+)	Physical Health	Itch – Activity & Clothing		Activity and clothing related quality of life impairment from itch (pruritis).
Adult (Ages 18+)	Physical Health	Itch – Mood & Sleep		Mood and sleep related quality of life impairment from itch (pruritis).

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Physical Health	Itch – Triggers		Assesses the subjective description of the triggers of itch with a checklist.
Adult (Ages 18+)	Physical Health	Dyspnea – Activity Motivation**		An adult’s relative dispositional tendency or preference toward being active or sedentary.
Adult (Ages 18+)	Physical Health	Dyspnea – Activity Requirements**		Impact of an adult's environment on their physical activity levels and external activity demands on an adult, such as work outside the home and aspects of the home environment (e.g., stairs).
Adult (Ages 18+)	Physical Health	Dyspnea – Airborne Exposure**		Environmental factors related to dyspnea, including exposure to airborne allergens, pollutants and smoke, and whether or not an adult is exposed to airborne factors in their environment that could impact their shortness of breath.
Adult (Ages 18+)	Physical Health	Dyspnea – Assistive Devices**		Availability and use of assistive devices (e.g., cane, grab bar, or oxygen) and whether or not an individual uses such devices.
Adult (Ages 18+)	Physical Health	Dyspnea – Characteristics**		Various descriptive aspects of a person’s experience of dyspnea, including quantitative and qualitative descriptions of the severity and intensity of shortness of breath as well as its frequency and duration.
Adult (Ages 18+)	Physical Health	Dyspnea – Emotional Response**		Emotions experienced related to dyspnea, including embarrassment, fear and worry, and how individuals may react emotionally to their shortness of breath.
Adult (Ages 18+)	Physical Health	Dyspnea – Functional Limitations		Impact of dyspnea (i.e. shortness of breath or difficulty breathing) on an adult’s ability to function while performing specific daily activities (e.g., dressing oneself without help, preparing meals, walking up 20 stairs).
Adult (Ages 18+)	Physical Health	Dyspnea – Severity		Severity of shortness of breath or difficulty breathing an adult experiences in response to various specific activities (the same activities assessed in Dyspnea Functional Limitations).
Adult (Ages 18+)	Physical Health	Dyspnea – Task Avoidance**		An adult’s decision to stop engaging in certain activities when the breathing discomfort associated with continuing the activity can no longer be tolerated.
Adult (Ages 18+)	Physical Health	Dyspnea – Time Extension**		Whether there has been a meaningful increase or decrease in the duration of time needed by an adult to perform a given task in the past 7 days (compared to 3 months ago) due to shortness of breath.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Physical Health	Gastrointestinal – Belly Pain		Severity of belly pain in adult populations . The intensity, nature (sharp vs. dull), frequency, bothersomeness, and predictability (e.g. ability to tell in advance when a pain episode would occur) all contribute towards belly pain severity.
Adult (Ages 18+)	Physical Health	Gastrointestinal – Constipation		Frequency and intensity of incomplete evacuation, rectal pain, straining, and hard stools, as well as the need for manual extraction of stool.
Adult (Ages 18+)	Physical Health	Gastrointestinal – Diarrhea		Frequency, form, bothersomeness, impact, controllability, and predictability of bowel urgency
Adult (Ages 18+)	Physical Health	Gastrointestinal – Disrupted Swallowing		Difficulty swallowing solid and soft foods, liquids, and pills as well as throat/chest pain and the sensation of food being stuck in the throat/chest.
Adult (Ages 18+)	Physical Health	Gastrointestinal – Gas and Bloating		Frequency and intensity/severity of bloating (i.e. feeling pressure or fullness), bloating appearance (i.e. belly swollen or larger than usual size), flatulence (i.e. passing gas), and abdominal sounds (i.e. gurgling or rumbling).
Adult (Ages 18+)	Physical Health	Gastrointestinal – Gastroesophageal Reflux		Symptoms associated with stomach contents leaking backwards from the stomach into the esophagus.
Adult (Ages 18+)	Physical Health	Gastrointestinal – Nausea and Vomiting		Frequency of vomiting, nausea, and poor appetite, as well as the predictability of nausea.
Adult (Ages 18+)	Physical Health	Gastrointestinal – Bowel Incontinence		Frequency of bowel incontinence, soiling, and gas incontinence (i.e. stool leakage while passing gas)
Adult (Ages 18+)	Physical Health	Physical Function		Self-reported capability rather than actual performance of physical activities. This includes the functioning of one’s upper extremities (dexterity), lower extremities (walking or mobility), and central regions (neck, back), as well as instrumental activities of daily living, such as running errands.
Adult (Ages 18+)	Physical Health	Physical Function	Mobility	Activities of physical mobility such as getting out of bed or a chair to activities such as running.
Adult (Ages 18+)	Physical Health	Physical Function	Upper Extremity	Activities that require use of the upper extremity including shoulder, arm, and hand activities. Examples include writing, using buttons, or opening containers.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Physical Health	Physical Function for Samples with Mobility Aid Users		For samples that may include individuals who use mobility aids such as wheelchairs.
Adult (Ages 18+)	Physical Health	Sleep Disturbance		Perceptions of sleep quality, sleep depth, and restoration associated with sleep.
Adult (Ages 18+)	Physical Health	Sleep-Related Impairment		Perceptions of alertness, sleepiness, and tiredness during usual waking hours, and the perceived functional impairments during wakefulness associated with sleep problems or impaired alertness.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Anal Discomfort with Sexual Activity (for Sexually Active People)	Anal irritation, pain, or bleeding during or after anal sex
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Bother Regarding Sexual Function - Female - Male	The extent to which people were bothered by aspects of sexual functioning
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Erectile Function (for Sexually Active Men)	Ability to achieve and maintain an erection for sexual activity.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Factors Interfering with Sexual Satisfaction	Perception of the degree to which various factors affect satisfaction with sex life. These factors include symptoms of disease and side effects from treatment and other issues that have been identified by participants.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Interest in Sexual Activity	Conscious awareness of wanting to engage in sexual activity.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Oral Discomfort with Sexual Activity (for Sexually Active People)	The degree of physical discomfort in the mouth, including pain and/or irritation, experienced with sexual activity

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Oral Dryness with Sexual Activity (for Sexually Active People)	The lack of saliva in the mouth experienced with sexual activity
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Orgasm – Ability (for Sexually Active People)	The degree to which the person has experienced a satisfying climax.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Orgasm - Pleasure (for Sexually Active People)	How pleasurable or satisfying the person's orgasms have felt.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Satisfaction with Sex Life	An overall evaluation of his or her sex life. No limitation is placed on what the person includes in his or her definition of “sex life.”
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Screeners	Asks about sex (gender), whether people are in a relationship that could involve sexual activity, and whether they have had any type of sexual activity with a partner in the past 30 days.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Sexual Activities - Female - Male	Frequency of engaging in specific intimate or sexual behaviors, either alone or with a partner.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Therapeutic Aids for Sexual Activity - Female - Male	Use of hormones, personal lubrications, medications, or devices intended to allow for or improve sexual function.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Vaginal Discomfort with Sexual Activity (for Sexually Active Women)	Physical discomfort of the vagina during and immediately following sexual activity.
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Vaginal Lubrication for Sexual Activity (for Sexually Active Women)	Wetness or dryness of the vagina during sexual activity.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Vulvar Discomfort with Sexual Activity – Clitoral (for Sexually Active Women)	The degree of physical discomfort, including pain, of the clitoris experienced with sexual activity
Adult (Ages 18+)	Physical Health	Sexual Function and Satisfaction	Vulvar Discomfort with Sexual Activity – Labial (for Sexually Active Women)	The degree of physical discomfort, including pain, of the labia experienced with sexual activity
Pediatric (Ages 8-17)	Physical Health	Pain – Behavior		Behaviors that typically indicate to others that an individual is experiencing pain. These actions or reactions can be verbal or nonverbal, and involuntary or deliberate.
Pediatric (Ages 8-17)	Physical Health	Pain – Interference		Consequences of pain on relevant aspects of one’s life. This includes the extent to which pain hinders engagement with social, cognitive, emotional, physical, and recreational activities.
Pediatric (Ages 8-17)	Physical Health	Pain Quality - Affective		Specific physical sensations and affective components associated with pain. Because pain can be felt and described in so many ways, this category of pain contains a variety of attributes, such as perceived temperature (e.g., cold), sensations (e.g., throbbing), and perceived affective qualities of pain (e.g., uncomfortable).
Pediatric (Ages 8-17)	Physical Health	Pain Quality - Sensory		Specific physical sensations and affective components associated with pain. Because pain can be felt and described in so many ways, this category of pain contains a variety of attributes, such as perceived temperature (e.g., cold), sensations (e.g., throbbing), and perceived affective qualities of pain (e.g., uncomfortable).
Pediatric (Ages 8-17)	Physical Health	Pain Quality		Specific physical sensations and affective components associated with pain. Because pain can be felt and described in so many ways, this category of pain contains a variety of attributes, such as perceived temperature (e.g., cold), sensations (e.g., throbbing), and perceived affective qualities of pain (e.g., uncomfortable).



Respondent	Category	Domain	Sub-Domain	Domain Definition
Pediatric (Ages 8-17)	Physical Health	Fatigue		Range of symptoms, from mild subjective feelings of tiredness to an overwhelming, debilitating, and sustained sense of exhaustion.
Pediatric (Ages 8-17)	Physical Health	Itch (PIQ-C)		Assesses itch (pruritus) symptoms and their impact on children's daily living. The PIQ-C includes items for children with skin conditions.
Pediatric (Ages 8-17)	Physical Health	Asthma Impact		Asthma-specific symptoms that include cough, wheeze, shortness of breath, and avoidance of triggers. Also, asthma-associated impacts such as missing school or activities with other children.
Pediatric (Ages 8-17)	Physical Health	Physical Stress Experience		The physically experienced sensations associated with responses to internal or external challenges including arousal, agitation, pain, and gastrointestinal distress.
Pediatric (Ages 8-17)	Physical Health	Physical Function	Mobility	Activities of physical mobility such as getting out of bed or a chair to activities such as running.
Pediatric (Ages 8-17)	Physical Health	Physical Function	Upper Extremity	Activities that require use of the upper extremity including shoulder, arm, and hand activities.
Pediatric (Ages 8-17)	Physical Health	Sleep-Related Disturbance		Assesses reported thoughts of one's sleep quality, and perceived difficulties with falling or staying asleep. Conceptual facets include sleep quality, sleep onset, and sleep continuity.
Pediatric (Ages 8-17)	Physical Health	Sleep-Related Impairment		Assesses perceptions of sleepiness during usual awake hours and reported impairments during the day associated with sleep problems or daytime sleepiness. Conceptual facets include daytime sleepiness, sleep offset, impact: cognitive, impact: activities, and impact: emotional.
Pediatric (Ages 8-17)	Physical Health	Physical Activity		Self-reported capability rather than actual performance of physical activities. This includes the functioning of one's upper extremities (dexterity), lower extremities (walking or mobility), and central regions (neck, back), as well as instrumental activities of daily living.
Pediatric (Ages 8-17)	Physical Health	Strength Impact		A child's capacity to perform functional activities of daily living that require significant amount of muscle force generation.
Parent Proxy (Ages 1-5)	Physical Health	Sleep-Related Disturbance		Sleep quality, sleep onset (difficulties falling asleep), sleep continuity (difficulties staying asleep).
Parent Proxy (Ages 1-5)	Physical Health	Sleep-Related Impairment		Perceptions of sleepiness during usual awake hours and reported impairments during the day associated with sleep problems or daytime sleepiness.



Respondent	Category	Domain	Sub-Domain	Domain Definition
Parent Proxy (Ages 5-17)	Physical Health	Pain – Behavior		Behaviors that typically indicate to others that an individual is experiencing pain. These actions or reactions can be verbal or nonverbal, and involuntary or deliberate.
Parent Proxy (Ages 5-17)	Physical Health	Pain – Interference		Consequences of pain on relevant aspects of one’s life. This includes the extent to which pain hinders engagement with social, cognitive, emotional, physical, and recreational activities.
Parent Proxy (Ages 5-17)	Physical Health	Fatigue		Range of symptoms, from mild subjective feelings of tiredness to an overwhelming, debilitating, and sustained sense of exhaustion.
Parent Proxy (Ages 5-17)	Physical Health	Itch		Assesses parents’ perception of itch (pruritus) symptoms and their impact on children’s daily living. The proxy PIQ-C includes items for parents’ perception of their children with skin conditions.
Parent Proxy (Ages 5-17)	Physical Health	Asthma Impact		Asthma-specific symptoms that include cough, wheeze, shortness of breath, and avoidance of triggers. Also, asthma-associated impacts such as missing school or activities with other children.
Parent Proxy (Ages 5-17)	Physical Health	Physical Stress Experience		The physically experienced sensations associated with responses to internal or external challenges including arousal, agitation, pain, and gastrointestinal distress.
Parent Proxy (Ages 5-17)	Physical Health	Physical Function	Mobility	Activities of physical mobility such as getting out of bed or a chair to activities such as running.
Parent Proxy (Ages 5-17)	Physical Health	Physical Function	Upper Extremity	Activities that require use of the upper extremity including shoulder, arm, and hand activities.
Parent Proxy (Ages 5-17)	Physical Health	Sleep-Related Disturbance		Sleep quality, sleep onset (difficulties falling asleep), sleep continuity (difficulties staying asleep).
Parent Proxy (Ages 5-17)	Physical Health	Sleep-Related Impairment		Perceptions of sleepiness during usual awake hours and reported impairments during the day associated with sleep problems or daytime sleepiness.
Parent Proxy (Ages 5-17)	Physical Health	Physical Activity		Self-reported capability rather than actual performance of physical activities. This includes the functioning of one’s upper extremities (dexterity), lower extremities (walking or mobility), and central regions (neck, back), as well as instrumental activities of daily living.
Parent Proxy (Ages 5-17)	Physical Health	Strength Impact		A child's capacity to perform functional activities of daily living that require significant amount of muscle force generation.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Mental Health				
Adult (Ages 18+)	Mental Health	Emotional Distress – Anxiety		Fear (fearfulness, panic), anxious misery (worry, dread), hyperarousal (tension, nervousness, restlessness), and somatic symptoms related to arousal (racing heart, dizziness).
Adult (Ages 18+)	Mental Health	Emotional Distress – Depression		Negative mood (sadness, guilt), views of self (self-criticism, worthlessness), and social cognition (loneliness, interpersonal alienation), as well as decreased positive affect and engagement (loss of interest, meaning, and purpose).
Adult (Ages 18+)	Mental Health	Emotional Distress – Anger		Angry mood (irritability, frustration), negative social cognitions (interpersonal sensitivity, envy, disagreeableness), and efforts to control anger.
Adult (Ages 18+)	Mental Health	Life Satisfaction		One's cognitive evaluation of life experiences and whether one likes his/her life or not.
Adult (Ages 18+)	Mental Health	Meaning and Purpose		A sense that life has purpose and there are good reasons for living, including hopefulness, optimism, goal-directedness, and feelings that one's life is worthy.
Adult (Ages 18+)	Mental Health	Positive Affect		Feelings that reflect a level of pleasurable engagement with the environment, such as happiness, joy, excitement, enthusiasm, and contentment.
Adult (Ages 18+)	Mental Health	Psychosocial Illness Impact – Negative		Direct negative psychosocial effect of cancer, distinct from general emotional distress.
Adult (Ages 18+)	Mental Health	Psychosocial Illness Impact – Positive		Positive psychosocial (emotional and social) outcomes of illness, previously conceptualized in various ways including post-traumatic growth, benefit-finding, and meaning making.
Adult (Ages 18+)	Mental Health	Alcohol – Alcohol Use		Drinking patterns, cue-based drinking, cravings to drink, and efforts to control drinking that indicate problematic drinking, particularly at the high end of the severity continuum.
Adult (Ages 18+)	Mental Health	Alcohol – Negative Consequences		Negative personal outcomes of alcohol use over the past 30 days. These items cover physical, mental, and social consequences of drinking.
Adult (Ages 18+)	Mental Health	Alcohol – Negative Expectancies		General attitudes about negative outcomes of alcohol use. These items cover physical, mental, and social negative expectancies of drinking.
Adult (Ages 18+)	Mental Health	Alcohol – Positive Consequences		Positive personal outcomes of alcohol use over the past 30 days. These items cover physical, mental, and social consequences of drinking.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Mental Health	Alcohol – Positive Expectancies		General attitudes about positive outcomes of alcohol use. These items cover physical, mental, and social positive expectancies of drinking.
Adult (Ages 18+)	Mental Health	Medication Adherence		Behaviors related to taking oral medication as prescribed by a doctor.
Adult (Ages 18+)	Mental Health	Smoking – Coping Expectancies All Smokers; Daily Smokers; Nondaily Smokers		Using smoking as a way to cope with various types of negative affect, affective consequences of not smoking, and the extent to which negative affect triggers smoking.
Adult (Ages 18+)	Mental Health	Smoking – Emotional/Sensory Expectancies All Smokers; Daily Smokers; Nondaily Smokers		Improved cognitive abilities (e.g., concentration), positive affective states (e.g., relaxation, contentment), and enjoyable sensorimotor sensations (e.g., from the ritual of lighting up a cigarette, smelling the cigarette, inhaling the smoke) that are experienced as a result of smoking.
Adult (Ages 18+)	Mental Health	Smoking – Negative Health Expectancies All Smokers; Daily Smokers; Nondaily Smokers		Health-related outcome expectancies of smoking (e.g., Smoking gives me a headache) and quitting (e.g., if I quit smoking I will breathe easier), and emotions such as worry (e.g., about developing serious health problems in the future).
Adult (Ages 18+)	Mental Health	Smoking – Negative Psychosocial Expectancies All Smokers; Daily Smokers; Nondaily Smokers		Friend and family disapproval (e.g., if I quit smoking my friends will respect me more), self-disapproval (e.g., I get upset when I think about my smoking), and internalization of social norms (e.g., I feel uncomfortable smoking around kids).
Adult (Ages 18+)	Mental Health	Smoking – Nicotine Dependence All Smokers; Daily Smokers; Nondaily Smokers		Tolerance, craving, withdrawal severity, temptations to smoke, and smoking as a behavioral priority.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Mental Health	Smoking – Social Motivations	All Smokers; Daily Smokers; Nondaily Smokers	Beliefs that: (a) smoking makes social situations more comfortable or enjoyable; (b) smoking provides a sense of camaraderie and belonging; (c) quitting smoking can negatively impact existing relationships with smokers; and (d) being in certain social situations increases smoking or the temptation to smoke.
Adult (Ages 18+)	Mental Health	Substance Use - Appeal	Past 30 Days; Past 3 months	Perceived positive aspects of substance use, including both increasing positive emotions (e.g., feeling happy and social) and alleviating negative emotions (e.g., reducing depression and anxiety). Requires a screening question to document the presence of some use: “In the past 30 days, have you used drugs other than alcohol or your prescribed medications?”
Adult (Ages 18+)	Mental Health	Substance Use - Prescription Pain Medication Misuse		Abuse of prescription pain medication. Requires a screening question to document the presence of a prescription for use: “In the past 3 months, did you have a prescription for pain medication?”
Adult (Ages 18+)	Mental Health	Substance Use - Severity	Past 30 Days; Past 3 months	Severity of substance use in adults. Requires a screening question to document the presence of some use: “In the past 3 months, have you used drugs other than alcohol or your prescribed medications?”
Adult (Ages 18+)	Mental Health	Cognitive Function		Mental acuity, concentration, verbal and nonverbal memory, verbal fluency, and perceived changes in these cognitive functions. The extent to which cognitive impairments interfere with daily functioning, whether other people observe cognitive impairments, and the impact of cognitive dysfunction on quality of life are also assessed.
Adult (Ages 18+)	Mental Health	Cognitive Function - Abilities		Patient-perceived functional abilities with regard to cognitive tasks, including the perception that one’s cognitive ability with regard to the domain of inquiry (e.g.concentration, memory) has not changed.
Adult (Ages 18+)	Mental Health	Self-Efficacy - General		Confidence in ability to deal effectively with a variety of stressful situations.
Adult (Ages 18+)	Mental Health	Self-Efficacy for Managing Chronic Conditions – Manage Emotions		Confidence to manage/control symptoms of anxiety, depression, helplessness, discouragement, frustration, disappointment and anger.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Mental Health	Self-Efficacy for Managing Chronic Conditions – Manage Daily Activities		Confidence in performing various activities of daily living (ADLs) without assistance. Items also assess exercise, sexual activities and managing activities in challenging situations (travelling, bad weather).
Adult (Ages 18+)	Mental Health	Self-Efficacy for Managing Chronic Conditions – Manage Meds/Treatment		Confidence in managing medication schedules of different complexity. Managing medication and other treatments in challenging situations such as when travelling, when running out of medication, and when adverse effects are encountered.
Adult (Ages 18+)	Mental Health	Self-Efficacy for Managing Chronic Conditions – Manage Social Interactions		Confidence in participating in social activities and getting help when necessary. Managing communication with others about their medical condition, including communication with health professionals.
Adult (Ages 18+)	Mental Health	Self-Efficacy for Managing Chronic Conditions – Manage Symptoms		Confidence to manage/control their symptoms, to manage their symptoms in different settings and to keep symptoms from interfering with work, sleep, relationships or recreational activities.
Adult (Ages 18+)	Mental Health	Healthcare Access Satisfaction		One's ability to get medical services, when needed (specific to adults with type 2 diabetes mellitus).
Adult (Ages 18+)	Mental Health	Illness Burden		The impact that a chronic medical condition(s) has on one's independent living, the ability to work (including working at home), social activities, and relationships.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Mental Health	Positive Attitudes Toward Complementary and Integrative Medicine		An individual's perception of the effectiveness of complementary and integrative medicine and the use of complementary and integrative medicine when seeking treatment. The items include level of preference for natural remedies and complementary and integrative medicine relative to conventional medicine.
Adult (Ages 18+)	Mental Health	Positive Outlook		Positive views of oneself, satisfaction with current life, and expectations for the future. The items capture self-worth, self-confidence, and levels of satisfaction with life circumstances as well as optimistic views of the future.
Adult (Ages 18+)	Mental Health	Positive Treatment Expectations		Patients' expectations about their treatment including the degree of confidence in treatment effectiveness, expectations of the benefit, perceived credibility of treatment concepts, and congruence of treatment with beliefs and attitudes about health.
Adult (Ages 18+)	Mental Health	Spirituality		Spiritual beliefs and experiences including receiving spiritual support, participating in religious activities and/or spiritual practices (e.g., meditation, prayer, yoga, visiting temples), and connection with a higher power.
Pediatric (Ages 8-17)	Mental Health	Emotional Distress – Anxiety		Fear (fearfulness, panic), anxious misery (worry, dread), hyperarousal (tension, nervousness, restlessness), and somatic symptoms related to arousal (racing heart, dizziness).
Pediatric (Ages 8-17)	Mental Health	Emotional Distress – Depressive Symptoms		Negative mood (sadness, guilt), views of self (self-criticism, worthlessness), and social cognition (loneliness, interpersonal alienation), as well as decreased positive affect and engagement (loss of interest, meaning, and purpose).
Pediatric (Ages 8-17)	Mental Health	Emotional Distress – Anger/Irritability		Angry mood (irritability, frustration), negative social cognitions (interpersonal sensitivity, envy, disagreeableness), and efforts to control anger.
Pediatric (Ages 8-17)	Mental Health	Life Satisfaction		Global and context-specific evaluations of a child's life. Conceptual facets include global evaluations of life, context-specific evaluations of life, assessments of life conditions, and comparisons of one's life with others' lives.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Pediatric (Ages 8-17)	Mental Health	Meaning and Purpose		A child's sense that life has purpose and there are good reasons for living, including hopefulness, optimism, goal-directedness, and feelings that one's life is worthy.
Pediatric (Ages 8-17)	Mental Health	Positive Affect		A child's momentary positive or rewarding affective experiences, such as feelings and mood associated with pleasure, joy, elation, contentment, pride, affection, happiness, engagement, and excitement.
Pediatric (Ages 8-17)	Mental Health	Psychological Stress Experiences		The thoughts or feelings about self and the world in the context of environmental or internal challenges. Items represent 3 facets of psychological stress reactions: feeling overwhelmed, perceived lack of control of capacity to manage one's life, and cognitive-perceptual disruption.
Pediatric (Ages 8-17)	Mental Health	Stigma	Stigma – Skin	A child's perceptions of self and publicly enacted negativity, prejudice, and discrimination as a result of disease-related manifestations. Banks include generic items for children with chronic conditions, as well as disease-specific items.
Pediatric (Ages 8-17)	Mental Health	Cognitive Function		Difficulties in cognitive abilities (e.g., memory, attention, and decision making), and difficulties in the application of such abilities to everyday tasks (e.g., planning, organizing, calculating, remembering, and learning).
Parent Proxy (Ages 1-5)	Mental Health	Emotional Distress – Anxiety		Fear (fearfulness, panic), anxious misery (worry, dread), hyperarousal (tension, nervousness, restlessness), social/separation anxiety (fear or distress when separating from caregivers), and somatic symptoms related to arousal (racing heart, dizziness).
Parent Proxy (Ages 1-5)	Mental Health	Emotional Distress – Depressive Symptoms		Negative mood (sadness, guilt), views of self (self-criticism, worthlessness), and social cognition (loneliness, interpersonal alienation); decreased positive affect, anhedonia (loss of interest, inability to engage in play), and engagement.
Parent Proxy (Ages 1-5)	Mental Health	Emotional Distress – Anger/Irritability		Angry mood (irritability, frustration, grouchiness), negative social cognitions (interpersonal sensitivity, envy, disagreeableness), behavior (tantrums), and efforts to control anger.
Parent Proxy (Ages 1-5)	Mental Health	Engagement - Curiosity		Young children's emotional, behavioral, and cognitive curiosity and interest, and initiative taking.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Parent Proxy (Ages 1-5)	Mental Health	Engagement - Persistence		Young children's sustained engagement and effort in problem solving and completing challenging activities and self-confidence
Parent Proxy (Ages 1-5)	Mental Health	Positive Affect		Momentary positive or rewarding affective experiences, such as feelings and mood associated with pleasure, joy, elation, contentment, pride, affection, happiness, engagement, and excitement.
Parent Proxy (Ages 1-5)	Mental Health	Self-Regulation – Flexibility		Young children's ability to adapt in response to environmental demands, changes, and expectations.
Parent Proxy (Ages 1-5)	Mental Health	Self-Regulation – Frustration Tolerance		Young children's recognition and regulation of emotions and behaviors in service of their goals (coping).
Parent Proxy (Ages 5-17)	Mental Health	Emotional Distress – Anxiety		Fear (fearfulness, panic), anxious misery (worry, dread), hyperarousal (tension, nervousness, restlessness), social/separation anxiety (fear or distress when separating from caregivers), and somatic symptoms related to arousal (racing heart, dizziness).
Parent Proxy (Ages 5-17)	Mental Health	Emotional Distress – Depressive Symptoms		Negative mood (sadness, guilt), views of self (self-criticism, worthlessness), and social cognition (loneliness, interpersonal alienation); decreased positive affect, anhedonia (loss of interest, inability to engage in play), and engagement.
Parent Proxy (Ages 5-17)	Mental Health	Emotional Distress – Anger/Irritability		Angry mood (irritability, frustration, grouchiness), negative social cognitions (interpersonal sensitivity, envy, disagreeableness), behavior (tantrums), and efforts to control anger.
Parent Proxy (Ages 5-17)	Mental Health	Life Satisfaction		Global and context-specific evaluations of a child's life. Conceptual facets include global evaluations of life, context-specific evaluations of life, assessments of life conditions, and comparisons of one's life with others' lives.
Parent Proxy (Ages 5-17)	Mental Health	Meaning and Purpose		A child's sense that life has purpose and there are good reasons for living, including hopefulness, optimism, goal-directedness, and feelings that one's life is worthy.
Parent Proxy (Ages 5-17)	Mental Health	Positive Affect		Momentary positive or rewarding affective experiences, such as feelings and mood associated with pleasure, joy, elation, contentment, pride, affection, happiness, engagement, and excitement.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Parent Proxy (Ages 5-17)	Mental Health	Psychological Stress Experiences		The thoughts or feelings about self and the world in the context of environmental or internal challenges. Items represent 3 facets of psychological stress reactions: feeling overwhelmed, perceived lack of control of capacity to manage one's life, and cognitive-perceptual disruption.
Parent Proxy (Ages 5-17)	Mental Health	Stigma Stigma – Skin		Parents' perceptions of self and publicly enacted negativity, prejudice, and discrimination as a result of disease-related manifestations. Banks include generic items for children with chronic conditions, as well as disease-specific items.
Parent Proxy (Ages 5-17)	Mental Health	Cognitive Function		Difficulties in cognitive abilities (e.g., memory, attention, and decision making), and difficulties in the application of such abilities to everyday tasks (e.g., planning, organizing, calculating, remembering, and learning).

Social Health

Adult (Ages 18+)	Social Health	Ability to Participate in Social Roles and Activities		Perceived ability to perform one's usual social roles and activities.
Adult (Ages 18+)	Social Health	Satisfaction with Participation in Discretionary Social Activities (v1.0)		Contentment with leisure interests and relationships with friends.
Adult (Ages 18+)	Social Health	Satisfaction with Participation in Social Roles (v1.0)		Satisfaction with performing one's usual social roles and activities
Adult (Ages 18+)	Social Health	Satisfaction with Social Roles and Activities (v2.0)		Satisfaction with performing one's usual social roles and activities (e.g., "I am satisfied with my ability to participate in family activities"). Two preliminary item banks were created based on PROMIS Wave 1 data (2005-2007): Satisfaction with Participation in Social Roles (v1.0) and Satisfaction with Participation in Discretionary Social Activities (v1.0). Supplemental data collection (2009-2010) with revised item pools led to the creation of one overall item bank: Satisfaction with Social Roles and Activities (v2.0).



Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Social Health	Social Isolation		Perceptions of being avoided, excluded, detached, disconnected from, or unknown by, others.
Adult (Ages 18+)	Social Health	Companionship		Perceived availability of someone with whom to share enjoyable social activities such as visiting, talking, celebrations, etc.
Adult (Ages 18+)	Social Health	Emotional Support		Perceived feelings of being cared for and valued as a person; having confident relationships.
Adult (Ages 18+)	Social Health	Informational Support		Perceived availability of helpful information or advice.
Adult (Ages 18+)	Social Health	Instrumental Support		Perceived availability of assistance with material, cognitive or task performance.
Pediatric (Ages 8-17)	Social Health	Peer Relationships		Quality of relationships with friends and other acquaintances.
Pediatric (Ages 8-17)	Social Health	Family Relationships		The subjective (affective, emotional, cognitive) experience of being involved with one's family, feeling like an important person in the family, of feeling accepted and cared for, and feeling that family members, especially parents, can be trusted and depended on for help and understanding.
Parent Proxy (Ages 1-5)	Social Health	Peer Relationships		Positive peer interactions, sociability (getting along well with others), and empathic behaviors.
Parent Proxy (Ages 1-5)	Social Health	Family Relationships		The subjective (affective, emotional, cognitive) experience of being involved with one's family, feeling like an important person in the family, of feeling accepted and cared for, and feeling that family members, especially parents, can be trusted and depended on for help and understanding.
Parent Proxy (Ages 5-17)	Social Health	Peer Relationships		Positive peer interactions, sociability (getting along well with others), and empathic behaviors.
Parent Proxy (Ages 5-17)	Social Health	Family Relationships		The subjective (affective, emotional, cognitive) experience of being involved with one's family, feeling like an important person in the family, of feeling accepted and cared for, and feeling that family members, especially parents, can be trusted and depended on for help and understanding.
Global Health				
Adult (Ages 18+)	Global	Global Health*		Overall evaluation of one's physical and mental health.



Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Global	Global Mental		Overall evaluation of one's mental health.
Adult (Ages 18+)	Global	Global Physical		Overall evaluation of one's physical health.
Pediatric (Ages 8-17)	Global	Global Health		Overall evaluation of one's physical and mental health.
Parent Proxy (Ages 5-17)	Global	Global Health		Overall evaluation of physical, mental health, and social health.
Profiles				
Adult (Ages 18+)	Profile	PROMIS-29 Profile (v2.1)		A collection of 4-item short forms assessing anxiety, depression, fatigue, pain interference, physical function, sleep disturbance, and ability to participate in social roles and activities as well as a single pain intensity item.
Adult (Ages 18+)	Profile	PROMIS-29+2 Profile v2.1 (PROPr)		Consists of the PROMIS 29+2 Profile v2.1 and two (2) PROMIS Cognitive Function Abilities items. This measure was created to incorporate the PROMIS Preference Scoring System.
Adult (Ages 18+)	Profile	PROMIS Profile CAT v1.0 – 29		A collection of 8 computer adaptive tests (CATs) assessing anxiety, depression, fatigue, pain interference, physical function, sleep disturbance, and ability to participate in social roles and activities as well as a single pain intensity item.
Adult (Ages 18+)	Profile	PROMIS-43 Profile (v2.1)		A collection of 6-item short forms assessing anxiety, depression, fatigue, pain interference, physical function, sleep disturbance, and ability to participate in social roles and activities as well as a single pain intensity item.
Adult (Ages 18+)	Profile	PROMIS-57 Profile (v2.1)		A collection of 8-item short forms assessing anxiety, depression, fatigue, pain interference, physical function, sleep disturbance, and ability to participate in social roles and activities as well as a single pain intensity item.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Brief Profile (Female)		A collection of brief (1-2) item short forms assessing participation and interest in sexual activity. Non-sexually active respondents are then asked about reasons for not participating in sexual activity, whereas sexually active respondents are then asked about orgasm, satisfaction with sex life, lubrication, and vaginal discomfort, vulvar discomfort.

Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Brief Profile (Male)		A collection of brief (1-2) item short forms assessing participation and interest in sexual activity. Non-sexually active respondents are then asked about reasons for not participating in sexual activity, whereas sexually active respondents are then asked about orgasm, satisfaction with sex life, and erectile function.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Brief Profile (Sexually Active Female)		A collection of brief (1-2) item short forms, intended for sexually active participants, that assesses interest in sexual activity, orgasm, satisfaction with sex life, lubrication, vaginal discomfort, and vulvar discomfort.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Brief Profile (Sexually Active Male)		A collection of brief (1-2) item short forms, intended for sexually active participants, that assesses interest in sexual activity, orgasm, satisfaction with sex life, and erectile function.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Female)		A collection of brief (1-4) item short forms assessing participation and interest in sexual activity. Non-sexually active respondents are then asked about reasons for not participating in sexual activity, whereas sexually active respondents are then asked about orgasm, satisfaction with sex life, lubrication, vaginal discomfort, vulvar discomfort, oral discomfort, oral dryness, and anal discomfort.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Male)		A collection of brief (1-4) item short forms assessing participation and interest in sexual activity. Non-sexually active respondents are then asked about reasons for not participating in sexual activity, whereas sexually active respondents are then asked about orgasm, satisfaction with sex life, erectile function, oral discomfort, oral dryness, and anal discomfort.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Non-Sexually Active Female)		A collection of brief (1-2) item short forms, intended for non-sexually active participants, that assesses interest and reasons for not participating in sexual activity.



Respondent	Category	Domain	Sub-Domain	Domain Definition
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Non-Sexually Active Male)		A collection of brief (1-2) item short forms, intended for non-sexually active participants, that assesses interest and reasons for not participating in sexual activity.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Sexually Active Female)		A collection of brief (1-4) item short forms, intended for sexually active participants, that assesses interest in sexual activity, orgasm, satisfaction with sex life, lubrication, vaginal discomfort, and vulvar discomfort.
Adult (Ages 18+)	Profile	Sexual Function and Satisfaction v2.0 Full Profile (Sexually Active Male)		A collection of brief (1-4) item short forms, intended for sexually active participants, that assesses interest in sexual activity, orgasm, satisfaction with sex life, and erectile function.
Pediatric (Ages 8-17)	Profile	PROMIS Pediatric-25 Profile		A collection of 4-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.
Pediatric (Ages 8-17)	Profile	PROMIS Pediatric-36 Profile		A collection of 5- and 6-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.
Pediatric (Ages 8-17)	Profile	PROMIS Pediatric-48 Profile		A collection of 7- and 8-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.
Parent Proxy (Ages 5-17)	Profile	PROMIS Parent Proxy-25 Profile		A collection of 4-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.
Parent Proxy (Ages 5-17)	Profile	PROMIS Parent Proxy-36 Profile		A collection of 5- and 6-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.
Parent Proxy (Ages 5-17)	Profile	PROMIS Parent Proxy-48 Profile		A collection of 7- and 8-item short forms assessing anxiety, depressive symptoms, fatigue, pain interference, physical function-mobility, and peer relationships, as well as a single pain intensity item.